



154

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07K 14/72	A1	(11) International Publication Number: WO 99/32519 (43) International Publication Date: 1 July 1999 (01.07.99)
<p>(21) International Application Number: PCT/SE98/02348</p> <p>(22) International Filing Date: 16 December 1998 (16.12.98)</p> <p>(30) Priority Data: 9704836-7 22 December 1997 (22.12.97) SE</p> <p>(71) Applicant (for all designated States except MG US): ASTRA PHARMA INC. [CA/CA]; 1004 Middlegate Road, Mississauga, Ontario L4Y 1M4 (CA).</p> <p>(71) Applicant (for MG only): ASTRA AKTIEBOLAG [SE/SE]; S-151 85 Södertälje (SE).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): AHMAD, Sultan [IN/CA]; Astra Research Centre Montreal, 7171 Fredrick-Banting, St. Laurent, Quebec H4S 1Z9 (CA); BANVILLE, Denis [CA/CA]; 595 Lajeunesse, Ste-Dorothee, Quebec H7X 3K4 (CA); FORTIN, Yves [CA/CA]; 2985 Avenue Douglas, Montreal, Quebec H3R 2E2 (CA); LEMBO, Paola [CA/CA]; Astra Research Centre Montreal, 7171 Fredrick-Banting, St. Laurent, Quebec H4S 1Z9 (CA); O'DONELL, Dajan [CA/CA]; Astra Research Centre Montreal, 7171 Fredrick-Banting, St. Laurent, Quebec H4S</p>	<p>1Z9 (CA). SHEN, Shi-Hsiang [CA/CA]; 161 Charwell Crescent, Beaconsfield, Quebec H9W 1C2 (CA).</p> <p>(74) Agent: ASTRA AKTIEBOLAG; Intellectual Property, S-151 85 Södertälje (SE).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p>	
<p>(54) Title: NOVEL G PROTEIN-COUPLED RECEPTOR</p> <p>(57) Abstract</p> <p>The present invention is directed to novel G protein-coupled receptors that are found predominantly in the dorsal root ganglia. The invention encompasses both receptor proteins as well as nucleic acids encoding the proteins. Angiotension 1 and III effects Calcium signalling in Cells transformed with DNA encoding the receptor. In addition, the present invention is directed to methods and compositions which utilize the receptors.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

NOVEL G PROTEIN-COUPLED RECEPTOR

Field of the Invention

The present invention is in the general field of biological receptors and the various uses
5 that can be made of such receptors. More specifically, the invention relates to nucleic acids
encoding novel G protein-coupled receptors and to the receptors per se.

Background and Prior Art

G protein-coupled receptors (GPCRs) constitute a family of proteins sharing a common
10 structural organization characterized by an extracellular N-terminal end, seven hydrophobic
alpha helices putatively constituting transmembrane domains and an intracellular C-
terminal domain. GPCRs bind a wide variety of ligands that trigger intracellular signals
through the activation of transducing G proteins (*Caron, et al., Rec. Prog. Horm. Res.*
48:277-290 (1993); *Freedman et al., Rec. Prog. Horm. Res. 51:319-353 (1996)*).

15

More than 300 GPCRs have been cloned thus far and it is generally assumed that there
exist well over 1000 such receptors. Mechanistically, approximately 50-60% of all
clinically relevant drugs act by modulating the functions of various GPCRs (*Cudermann, et*
al., J. Mol. Med. 73:51-63 (1995)). Of particular interest are receptors located in dorsal root
20 ganglia. This region of the central nervous system is densely innervated with primary or
afferent sensory neurons involved in the transmission, modulation and sensation of pain.
Thus, receptors from this region may be used in assays for the identification of new agents
for anesthesia and analgesia

25 Summary of the Invention

The present invention is based upon the discovery of a novel G protein-coupled receptor
which is distinct from previously reported receptors in terms of structure and in being
expressed preferentially in dorsal root ganglia. One dorsal root receptor (DRR) has been
isolated and sequenced from the rat and six from the human. The rat receptor was given the

designation rDRR-1 and its amino acid sequence is shown as SEQ ID NO:1. The human receptors were designated as

hDRR-1 (SEQ ID NO:3);

hDRR-2 (SEQ ID NO:5);

5 hDRR-3 (SEQ ID NO:7):

hDRR-4 (SEQ ID NO:9);

hDRR-5 (SEQ ID NO:11); and

hDRR-6 (SEQ ID NO:13).

Unless otherwise specified, the term "DRR" as used herein refers to all of the receptors
10 from both human and rat .

In its first aspect, the invention is directed to proteins, except as existing in nature, comprising the amino acid sequence consisting functionally of a rat or human DRR as shown in SEQ ID NO:1, 3, 5, 7, 9, 11, or 13. The term "consisting functionally of" is
15 intended to include any receptor protein whose sequence has undergone additions, deletions or substitutions which do not substantially alter the functional characteristics of the receptor. Thus, the invention encompasses proteins having exactly the same amino acid sequence as shown in the sequence listing, as well as proteins with differences that are not substantial as evidenced by their retaining the basic, qualitative binding properties of
20 the DRR receptor. The invention further encompasses substantially pure proteins consisting essentially of a DRR amino acid sequence, antibodies that bind specifically to a DRR (i.e. that have at least a 100 fold greater affinity for the DRR than any other naturally occurring non-DRR protein), and antibodies made by a process involving the injection of pharmaceutically acceptable preparations of such proteins into an animal capable of
25 antibody production. In a preferred embodiment, monoclonal antibody to human or rat DRR is produced by injecting a pharmaceutically acceptable preparation of the receptor into a mouse and then fusing mouse spleen cells with myeloma cells.

The invention is also directed to a substantially pure polynucleotide encoding a protein
30 comprising the amino acid sequence consisting functionally of the sequence of rat DRR (as

shown in SEQ ID NO:1) or a human DRR (as shown in SEQ ID NOs 3, 5, 7, 9, 11 or 13). This aspect of the invention encompasses polynucleotides encoding proteins consisting essentially of the amino acid sequences shown in the sequence listing, expression vectors comprising such polynucleotides, and host cells transformed with such vectors. Also
5 included are the recombinant rat and human DRR proteins produced by host cells made in this manner.

Preferably, the polynucleotide encoding rat DRR has the nucleotide sequence shown in SEQ ID NO:2 and the polynucleotide encoding a human DRR has the nucleotide sequence
10 shown in SEQ ID NO: 3, 5, 7, 9, 11 or 13. It is also preferred that the vectors and host cells used for the expression of DRR contain these particular polynucleotides.

In another aspect, the present invention is directed to a method for assaying a test compound for its ability to bind to a rat or human DRR. The method is performed by
15 incubating a source of DRR with a ligand known to bind to the receptor and with the test compound. The source of the DRR should be substantially free of other types of G protein-coupled receptors, i.e. greater than 85% of such receptors present should correspond to the DRR. Upon completion of incubation, the ability of the test compound to bind to the DRR is determined by the extent to which ligand binding has been displaced. The rat DRR
20 should, preferably correspond to rDDR-1 as shown in SEQ ID NO:1. The human receptor should preferably be hDDR-1 (SEQ ID NO:3); hDDR-2 (SEQ ID NO:5); hDDR-3 (SEQ ID NO:7); hDDR-4 (SEQ ID NO:9); hDDR-5 (SEQ ID NO:11); or hDDR-6 (SEQ ID NO:13). Either transformed cells expressing recombinant DRR may be used in the assays or membranes can be prepared from the cells and used. Although not essential, the assay can
25 be accompanied by the determination of the activation of a second messenger pathway such as the adenylyl cyclase pathway. This should help to determine whether a compound that binds to DRR is acting as an agonist or antagonist.

An alternative method for determining if a test compound is an agonist of any of the
30 DRRs disclosed herein is to use a cell signaling assay, e.g., an assay measuring either

intracellular adenylyl cyclase activity or intracellular calcium concentration. The test compound is incubated with cells expressing the DRR but substantially free of other G protein-coupled receptors, typically a cell transfected with an expression vector encoding the DRR. Test compounds that are agonists are identified by their causing a statistically significant change in the results obtained from the cell signaling assay when compared to control transfectants not exposed to test compound. For example, the cells exposed to the test compound may show a significant increase in adenylyl cyclase activity or in intracellular calcium concentration.

- 10 The invention also encompasses a method for determining if a test compound is an antagonist of a DRR which relies upon the known activation of G protein-coupled receptors that occurs when such receptors are expressed in large amounts. This method requires that DNA encoding the receptor be incorporated into an expression vector so that it is operably linked to a promoter and that the vector then be used to transfect an
- 15 appropriate host. In order to produce sufficient receptor to result in constitutive receptor activation (i.e., activation in the absence of natural ligand), expression systems capable of copious protein production are preferred, e.g., the DRR DNA may be operably linked to a CMV promoter and expressed in COS or HEK293 cells. After transfection, cells with activated receptors are selected based upon their showing increased activity in a cell
- 20 signaling assay relative to comparable cells that have either not been transfected or that have been transfected with a vector that is incapable of expressing functional DRR. Typically, cells will be selected either because they show a statistically significant increase in intracellular adenylyl cyclase activity or a statistically significant increase in intracellular calcium concentration. The selected cells are contacted with the test
- 25 compound and the cell signaling assay is repeated to determine if this results in a decrease in activity relative to control cells not contacted with the test compound. For example, a statistically significant decrease in either adenylyl cyclase activity or calcium concentration relative to control cells would indicate that the test compound is an antagonist of the DRR. Any of the DRRs disclosed herein may be used in these assays.

- Assays for compounds interacting with a DRR may be performed by incubating a source containing the DRR but substantially free of other G protein-coupled receptors (e.g. a stably transformed cell) with angiotensin II or III in both the presence and absence of test compound and measuring the modulation of intracellular calcium concentration. A
- 5 significant increase or decrease in angiotensin-stimulated calcium displacement in response to test compound is indicative of an interaction occurring at the DRR. The receptors that may be used in these assays include rat DRR-1 and human DRR-1, DRR-2, DRR-3, DRR-4, DRR-5 and DRR-6.
- 10 In another aspect, the present invention is directed to a method for assaying a test compound for its ability to alter the expression of a rat or human DRR. This method is performed by growing cells expressing the DRR, but substantially free of other G protein-coupled receptors, in the presence of the test compound. Cells are then collected and the expression of the DRR is compared with expression in control cells grown under
- 15 essentially identical conditions but in the absence of the test compound. The rat receptor is preferably rDRR-1 and the human receptor may be DRR-1; DRR-2; DRR-3; DRR-4; DRR-5; or DRR-6.
- A preferred test compound is an oligonucleotide at least 15 nucleotides in length
- 20 comprising a sequence complimentary to the sequence of the DRR used in the assay.

Brief Description of the Drawings

Figure 1. Nucleotide sequence of rDDR-1: Clone 3B-32, encoding rDDR-1, was isolated from a rat genomic library using the Promoter Finder Walking Kit (see Methods, Clontech).

5

The cloned gene was deposited with the international depositary authority Deutsche Sammlung Von Mikroorganismen Und Zellkulturen GmbH at the address Mascheroder Weg 1 B, D-3300 Braunschweig, Germany. The deposit was made on November 27, 1997 and was given the accession number DSM 11877.

10

Figure 2. Deduced amino acid sequence of DDR-1: Clone 3B-32 codes for a 337 amino acid protein. The amino acid sequence begins with the first ATG in the nucleotide sequence.

15 Figure 3. Alignment of the deduced amino acid sequences of clone 3B-32 (rDDR-1) with its five most homologous sequences. The boxed and shaded residues are the ones that are identical to the rDDR-1 sequence.

Figure 4. Amino acid alignment of the human DDR homologs: The amino acid sequence
20 of all 6 human homologs of rDDR-1 (hDDR-1; hDDR-2; hDDR-3; hDDR-4; hDDR-5; and hDDR-6) are aligned. The amino acid residues differing from the clone 36 (HUMAN36.PR) are boxed. The degree of identity among these sequences ranges from 77% to almost 100%.

25 Definitions

The description that follows uses a number of terms that refer to recombinant DNA technology. In order to provide a clear and consistent understanding of the specification and claims, including the scope to be given such terms, the following definitions are provided.

30

Cloning vector: A plasmid or phage DNA or other DNA sequence which is able to replicate autonomously in a host cell, and which is characterized by one or a small number of restriction endonuclease recognition sites. A foreign DNA fragment may be spliced into the vector at these sites in order to bring about the replication and cloning of the fragment.

- 5 The vector may contain a marker suitable for use in the identification of transformed cells. For example, markers may provide tetracycline resistance or ampicillin resistance.

Expression vector: A vector similar to a cloning vector but which is capable of inducing the expression of the DNA that has been cloned into it, after transformation into a host.

- 10 The cloned DNA is usually placed under the control of (i.e., operably linked to) certain regulatory sequences such as promoters or enhancers. Promoter sequences may be constitutive, inducible or repressible.

- Substantially pure:** As used herein, "substantially pure" means that the desired product is essentially free from contaminating cellular components. A "substantially pure" protein or nucleic acid will typically comprise at least 85% of a sample, with greater percentages being preferred. Contaminants may include proteins, carbohydrates or lipids. One method for determining the purity of a protein or nucleic acid is by electrophoresing a preparation in a matrix such as polyacrylamide or agarose. Purity is evidenced by the appearance of a single band after staining. Other methods for assessing purity include chromatography and analytical centrifugation.
- 15
20

- Host:** Any prokaryotic or eukaryotic cell that is the recipient of a replicable expression vector or cloning vector is the "host" for that vector. The term encompasses prokaryotic or eukaryotic cells that have been engineered to incorporate a desired gene on its chromosome or in its genome. Examples of cells that can serve as hosts are well known in the art, as are techniques for cellular transformation (see e.g. Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd ed. Cold Spring Harbor (1989)).
- 25

- 30 **Promoter:** A DNA sequence typically found in the 5' region of a gene, located proximal

to the start codon. Transcription is initiated at the promoter. If the promoter is of the inducible type, then the rate of transcription increases in response to an inducing agent.

Complementary Nucleotide Sequence: A complementary nucleotide sequence, as used
5 herein, refers to the sequence that would arise by normal base pairing. For example, the nucleotide sequence 5'-AGAC-3' would have the complementary sequence 5'-GTCT-3'.

Expression: Expression is the process by which a polypeptide is produced from DNA. The process involves the transcription of the gene into mRNA and the translation of this
10 mRNA into a polypeptide.

Detailed Description of the Invention

The present invention is directed to DRR receptor proteins, genetic sequences coding for
15 the receptors, a method for assaying compounds for binding to DRR receptors and a method for assaying compounds for their ability to alter DRR expression. The receptors and their nucleic acids are defined by their structures (as shown in figures 1, 2 and 4; and SEQ ID numbers 1-14).

20 It will be understood that the present invention encompasses not only sequences identical to those shown in the figures and sequence listing, but also sequences that are essentially the same and sequences that are otherwise substantially the same and which result in a receptor retaining the basic binding characteristics of the DRR. For example, it is well known that techniques such as site-directed mutagenesis may be used to introduce
25 variations in a protein's structure. Variations in a DRR protein introduced by this or some similar method are encompassed by the invention provided that the resulting receptor retains the basic qualitative binding characteristics of the unaltered DRR. Thus, the invention relates to proteins comprising amino acid sequences consisting functionally of the sequence of SEQ ID NO:1 (rat) and SEQ ID numbers 3, 5, 7, 9, 11 and 14 (human).

30

I. Nucleic Acid Sequences Coding for DRR

DNA sequences coding for DRRs are expressed exclusively, or at least highly preferentially, in dorsal root ganglia and these ganglia may serve as a source for the isolation of nucleic acids coding for the receptors. In addition, cells and cell lines that
5 express a rat or human DRR may serve as a source for nucleic acid. These may either be cultured cells that have not undergone transformation or cell lines specifically engineered to express recombinant DRR.

In all cases, poly A⁺ mRNA is isolated from the dorsal root ganglia, reverse transcribed
10 and cloned. The cDNA library thus formed may then be screened using probes derived from the sequences shown in the accompanying sequence listing as SEQ ID number 2, 4, 6, 8, 10, 12 or 14, depending upon the particular DRR being isolated. Probes should typically be at least 14 bases in length and should be derived from a portion of the DRR sequence that is poorly conserved (see Figures 3 and 4). Screening can also be performed using
15 genomic libraries with one DRR gene, or a portion of the gene, serving as a probe in the isolation of other DRR genes. For example, full length rDRR-1 may be labeled and used to screen a human genomic library for the isolation of hDRR-1, hDRR-2 etc. (see Examples section).

20 Alternatively genomic DNA libraries can be used to isolate DRR genes by performing PCR amplifications with primers located at either end of genes (see Examples section for a description of procedures). For example, human genomic DNA may be amplified using the primers:

25 5'-GCAAGCTTTCTGAGCATGGATCCAACCGTC, and
5'-CCCTCAGATCTCCAATTTGCTTCCCGACAG.

This will serve to amplify all six of the human DRR genes identified herein as hDRR-1; hDRR-2; hDRR-3; hDRR-4; hDRR-5; and hDRR-6. These may then be cloned into an appropriate vector, e.g. pGEM-T (Promega), for DNA sequence analysis.

5 II. Antibodies to Rat and Human DRRs

The present invention is also directed to antibodies that bind specifically to a rat or human DRR and to a process for producing such antibodies. Antibodies that "bind specifically to a DRR" are defined as those that have at least a one hundred fold greater affinity for the DRR than for any other protein. The process for producing such antibodies may involve
10 either injecting the DRR protein itself into an appropriate animal or, preferably, injecting short peptides made to correspond to different regions of the DRR. The peptides should be at least five amino acids in length and should be selected from regions believed to be unique to the particular DRR protein being targeted. Thus, highly conserved transmembrane regions should generally be avoided in selecting peptides for the generation
15 of antibodies. Methods for making and detecting antibodies are well known to those of skill in the art as evidenced by standard reference works such as: (*Harlow et al., Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory, N.Y. (1988)*); *Klein, Immunology: The Science of Self-Nonself Discrimination (1982)*; *Kennett, et al., Monoclonal Antibodies and Hybridomas: A New Dimension in Biological Analyses (1980)*;
20 and *Campbell, "Monoclonal Antibody Technology," in Laboratory Techniques in Biochemistry and Molecular Biology, (1984)*).

"Antibody," as used herein, is meant to include intact molecules as well as fragments which retain their ability to bind to antigen (e.g., Fab and F(ab)₂ fragments). These fragments are
25 typically produced by proteolytically cleaving intact antibodies using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab)₂ fragments). The term "antibody" also refers to both monoclonal antibodies and polyclonal antibodies. Polyclonal antibodies are derived from the sera of animals immunized with the antigen. Monoclonal antibodies can be prepared using hybridoma technology (*Kohler, et al., Nature 256:495*
30 (1975); *Hammerling, et al., in: Monoclonal Antibodies and T-Cell Hybridomas, Elsevier,*

M.Y., pp. 563-681 (1981)). In general, this technology involves immunizing an animal, usually a mouse, with either intact DRR or a fragment derived from the DRR. The splenocytes of the immunized animals are extracted and fused with suitable myeloma cells, e.g., SP2O cells. After fusion, the resulting hybridoma cells are selectively maintained in
5 HAT medium and then cloned by limiting dilution (Wands, et al., *Gastroenterology* 80:225-232 (1981)). The cells obtained through such selection are then assayed to identify clones which secrete antibodies capable of binding to the DRR.

The antibodies, or fragments of antibodies, of the present invention may be used to detect
10 the presence of DRR protein using any of a variety of immunoassays. For example, the antibodies may be used in radioimmunoassays or in immunometric assays, also known as "two-site" or "sandwich" assays (see Chard, T., *"An Introduction to Radioimmune Assay and Related Techniques," in Laboratory Techniques in Biochemistry and Molecular Biology, North Holland Publishing Co., N.Y. (1978)*). In a typical immunometric assay, a
15 quantity of unlabeled antibody is bound to a solid support that is insoluble in the fluid being tested, e.g., blood, lymph, cellular extracts, etc. After the initial binding of antigen to immobilized antibody, a quantity of detectably labeled second antibody (which may or may not be the same as the first) is added to permit detection and/or quantitation of bound antigen (see e.g. *Radioimmune Assay Method*, Kirkham et al., ed., pp. 199-206. E & S.
20 Livingstone, Edinburgh (1970)). Many variations of these types of assays are known in the art and may be employed for the detection of the DRR.

Antibodies to a rat or human DRR may also be used in the purification of either the intact receptor or fragments of the receptor (see generally, *Dean et al., Affinity Chromatography, A Practical Approach, IRL Press (1986)*). Typically, antibody is immobilized on a
25 chromatographic matrix such as Sepharose 4B. The matrix is then packed into a column and the preparation containing the DRR desired is passed through under conditions that promote binding, e.g., under conditions of low salt. The column is then washed and bound DRR is eluted using a buffer that promotes dissociation from antibody, e.g., buffer having

an altered pH or salt concentration. The eluted DRR may be transferred into a buffer of choice, e.g., by dialysis, and either stored or used directly.

III. Radioligand Assay for Receptor Binding

5 One of the main uses for DRR nucleic acids and recombinant proteins is in assays designed to identify agents capable of binding to DRR receptors. Such agents may either be agonists, mimicking the normal effects of receptor binding, or antagonists, inhibiting the normal effects of receptor binding. Of particular interest is the identification of agents which bind to the DRR and modulate adenylyl cyclase activity in the cells. These agents have
10 potential therapeutic application as either analgesics or anesthetics.

In radioligand binding assays, a source of DRR is incubated together with a ligand known to bind to the receptor and with the compound being tested for binding activity. The preferred source for DRR is cells, preferably mammalian cells, transformed to recombinantly express the receptor. The cells selected should not express a substantial
15 amount of any other G protein-coupled receptors that might bind to ligand and distort results. This can easily be determined by performing binding assays on cells derived from the same tissue or cell line as those recombinantly expressing DRR but which have not undergone transformation.

20 The assay may be performed either with intact cells or with membranes prepared from the cells (*see e.g. Wang, et al., Proc. Natl. Acad. Sci. U.S.A. 90:10230-10234 (1993)*). The membranes are incubated with a ligand specific for the DRR receptor and with a preparation of the compound being tested. After binding is complete, receptor is separated from the solution containing ligand and test compound, e.g. by filtration, and the amount of
25 binding that has occurred is determined. Preferably, the ligand used is detectably labeled with a radioisotope such as ¹²⁵I. However, if desired, fluorescent or chemiluminescent labels can be used instead. Among the most commonly used fluorescent labeling compounds are fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthalaldehyde and fluorescamine. Useful chemiluminescent compounds
30 include luminol, isoluminol, theromatic acridinium ester, imidazole, acridinium salt, and

oxalate ester. Any of these agents which can be used to produce a ligand suitable for use in the assay.

Nonspecific binding may be determined by carrying out the binding reaction in the presence of a large excess of unlabeled ligand. For example, labeled ligand may be incubated with receptor and test compound in the presence of a thousandfold excess of unlabeled ligand. Nonspecific binding should be subtracted from total binding, i.e. binding in the absence of unlabeled ligand, to arrive at the specific binding for each sample tested. Other steps such as washing, stirring, shaking, filtering and the like may be included in the assays as necessary. Typically, wash steps are included after the separation of membrane-bound ligand from ligand remaining in solution and prior to quantitation of the amount of ligand bound, e.g., by counting radioactive isotope. The specific binding obtained in the presence of test compound is compared with that obtained in the presence of labeled ligand alone to determine the extent to which the test compound has displaced receptor binding.

15

In performing binding assays, care must be taken to avoid artifacts which may make it appear that a test compound is interacting with the DRR receptor when, in fact, binding is being inhibited by some other mechanism. For example, the compound being tested should be in a buffer which does not itself substantially inhibit the binding of ligand to DRR and should, preferably, be tested at several different concentrations. Preparations of test compound should also be examined for proteolytic activity and it is desirable that antiproteases be included in assays. Finally, it is highly desirable that compounds identified as displacing the binding of ligand to DRR receptor be reexamined in a concentration range sufficient to perform a Scatchard analysis on the results. This type of analysis is well known in the art and can be used for determining the affinity of a test compounds for receptor (see e.g., Ausubel, et al., *Current Protocols in Molecular Biology*, 11.2.1-11.2.19 (1993); *Laboratory Techniques and Biochemistry and Molecular Biology*, Work, et al., ed., N.Y. (1978) etc.). Computer programs may be used to help in the analysis of results (see e.g., Munson, P., *Methods Enzymol.* 92:543-577 (1983); McPherson, G.A., *Kinetic, EBDA*

25

Ligand. Lowry-A Collection of Radioligand Binding Analysis Programs. Elsevier-Biosoft, U.K. (1985)).

The activation of receptor by the binding of ligand may be monitored using a number of
5 different assays. For example, adenylyl cyclase assays may be performed by growing cells in
wells of a microtiter plate and then incubating the various wells in the presence or absence
of test compound. cAMP may then be extracted in ethanol, lyophilized and resuspended in
assay buffer. Assay of cAMP thus recovered may be carried out using any method for
determining cAMP concentration, e.g. the Biotrack cAMP Enzyme-immunoassay System
10 (Amersham) or the Cyclic AMP [3H] Assay System (Amersham). Typically, adenylyl
cyclase assays will be performed separately from binding assays, but it may also be
possible to perform binding and adenylyl cyclase assays on a single preparation of cells.
Other "cell signaling assays" that can be used to monitor receptor activity are described
below.

15

IV. Identification of DRR Agonists and Antagonists Using Cell Signaling Assays

DRRs may also be used to screen for drug candidates using cell signaling assays. To
identify DRR agonists, the DNA encoding a receptor is incorporated into an expression
vector and then transfected into an appropriate host. The transformed cells are then
20 contacted with a series of test compounds and the effect of each is monitored. Among the
assays that can be used are assays measuring cAMP production (see discussion above),
assays measuring the activation of reporter gene activity, or assays measuring the
modulation of the binding of GTP-gamma-S.

25 Cell signaling assays may also be used to identify DRR antagonists. G protein-coupled
receptors can be put in their active state even in the absence of their cognate ligand by
expressing them at very high concentration in a heterologous system. For example, receptor
may be overexpressed using the baculovirus infection of insect Sf9 cells or a DRR gene
may be operably linked to a CMV promoter and expressed in COS or HEK293 cells. In this
30 activated constitutive state, antagonists of the receptor can be identified in the absence of

ligand by measuring the ability of a test compound to inhibit constitutive cell signaling activity. Appropriate assays for this are, again, cAMP assays, reporter gene activation assays or assays measuring the binding of GTP-gamma-S.

- 5 One preferred cell signaling assay is based upon the observation that cells stably transfected with DRRs show a change in intracellular calcium levels in response to incubation in the presence of angiotensin II or III (see Example 5). Thus, a procedure can be used to identify DRR agonists or antagonists that is similar to the radioreceptor assays discussed above except that angiotensin II or III is used instead of a labeled ligand and
- 10 calcium concentration is measured instead of bound radioactivity. The concentration of calcium in the presence of test compound and angiotensin II or III is compared with that in the presence of angiotensin II or III alone to determine whether the test compound is interacting at the DRR receptor. A statistically significant increase in intracellular calcium in response to test compound indicates that the test compound is acting as an agonist
- 15 whereas a statistically significant decrease in intracellular calcium indicates that it is acting as an antagonist.

V. Assay for Ability to Modulate DRR Expression

- One way to either increase or decrease the biological effects of a DRR is to alter the
- 20 extent to which the receptor is expressed in cells. Therefore, assays for the identification of compounds that either inhibit or enhance expression are of considerable interest. These assays are carried out by growing cells expressing a DRR in the presence of a test compound and then comparing receptor expression in these cells with expression in cells grown under essentially identical conditions but in the absence of the test compound. As in
- 25 the binding assays discussed above, it is desirable that the cells used be substantially free of competing G protein-coupled receptors. One way to quantitate receptor expression is to fuse the DRR sequence to a sequence encoding a peptide or protein that can be readily quantitated. For example, the DRR sequence may be ligated to a sequence encoding haemagglutinin as described in Example 5 and used to stably transfect cells. After

incubation with test compound the hemagglutinin/receptor complex can be immunoprecipitated and western blotted with anti- haemagglutinin antibody. Alternatively, Scatchard analysis of binding assays may be performed with labeled ligand to determine receptor number. The binding assays may be carried out as discussed above
5 and will preferably utilize cells that have been engineered to recombinantly express DRR.

A preferred group of test compounds for inclusion in the DRR expression assay consists of oligonucleotides complementary to various segments of the DRR nucleic acid sequence. These oligonucleotides should be at least 15 bases in length and should be derived from
10 non-conserved regions of the receptor nucleic acid sequence. Sequences may be based upon those shown as SEQ ID numbers 2, 4, 6, 8, 10, 12 or 14.

Oligonucleotides which are found to reduce receptor expression may be derivatized or conjugated in order to increase their effectiveness. For example, nucleoside
15 phosphorothioates may be substituted for their natural counterparts (see Cohen, J., Oligodeoxynucleotides, Antisense Inhibitors of Gene Expression, CRC Press (1989)). The oligonucleotides may be delivered to a patient in vivo for the purpose of inhibiting DRR expression. When this is done, it is preferred that the oligonucleotide be administered in a form that enhances its uptake by cells. For example, the oligonucleotide may be delivered
20 by means of a liposome or conjugated to a peptide that is ingested by cells (see e.g., U.S. Patent Nos. 4,897,355 and 4,394,448; see also non-U.S. patent documents WO 8903849 and EP 0263740). Other methods for enhancing the efficiency of oligonucleotide delivery are well known in the art and are also compatible with the present invention.

25 Having now described the invention, the same will be more readily understood through reference to the following Examples which are provided by way of illustration and which are not intended to limit the scope of the invention.

EXAMPLES**Example 1: Cloning of Rat DRR-1**

Isolation of cDNA fragment.

5

Degenerate oligonucleotides were synthesized to highly conserved regions of G-protein coupled receptors (transmembrane spanning domains 2 and 7) with the following nucleotide sequences:

10 5' GG CCG TCG ACT TCA TCG TC(A/T) (A/C)(T/C)C TI(G/T) CI(T/C) TIG
C(A/C/G/T)G 3' (TM2:sense) SEQ ID NO:15; and

5' (A/G)(C/A/T)(A/T) (A/G)CA (A/G)TA IAT IAT IGG (A/G)TT 3'
(TM7:antisense) SEQ ID NO:16.

15

Poly A+ mRNA was isolated from cultured fetal rat dorsal root ganglia (Sprague-Dawley). The mRNA was reverse transcribed using the First Strand cDNA Synthesis kit (Pharmacia Biotech), subjected to an amplification reaction by polymerase chain reaction (PCR) using Ampli-Taq DNA (Perkin-Elmer Cetus) polymerase under the following conditions: 3 minutes at 94 °C, 40 cycles of 1 minute at 94 °C, 45°C and 72 °C. A cDNA PCR fragment corresponding to approximately 650 bps was isolated and subcloned in pGEM-T-vector (Promega Corporation). The nucleotide sequence of the recombinant clone was determined using the T7-dideoxy chain termination sequencing kit (Pharmacia Biotech) and was found to be unique based upon searches of Genbank/EMBL
25 databases.

The full length rat DRR-1 sequence was obtained from rat genomic DNA using the 650 base pair fragment and the "Promoter Finder DNA Walking kit" (Clontech, cat # K1806-1). This kit contains five libraries of uncloned, adaptor-ligated genomic DNA fragments. The
30 procedure involves two consecutive PCR reactions. Both reactions were done using the

"Advantage Tth Polymerase Mix" also obtained from Clontech, following the conditions recommended by the vendor. The first PCR reaction was performed with the outer adaptor primer (AP1) provided in the kit and an outer, gene-specific primer (GSP1) derived from the sequence of the DRR-1 PCR fragment. The primary PCR mixtures were diluted and
5 used as a template for the secondary (nested) PCR reaction with the nested adapter primer (AP2) and a nested gene specific primer (GSP2). To obtain the sequence of the rat DRR-1 gene upstream of the sequence of the original PCR fragment, the following oligonucleotides were used:

- 10 GSP1: oligo YF3B59-B, 5'-CGCAGATGAGGTAGTACAGCATCAC SEQ ID NO:17
GSP2: oligo MML-R1, 5'-CTGTGAGAGAGATGGTAACATACAG SEQ ID NO:18

From the first library, a fragment AP2-MMLR1 of 1.9 Kb was obtained and from the third library, a fragment of approximately 1.0 Kb was obtained. To identify the sequence
15 downstream of the known sequence, the following primers were used:

- GSP1: oligo YF3B59-F2, 5'-GCATCCTTGACTGGTTCTTCTCAG SEQ ID NO:19
GSP2: oligo MML-F1, 5'-GGGTGAGACTCATCATCATTTGTGG. SEQ ID NO:20

- 20 A fragment MMLF1-AP2 of approximately 1 Kb was obtained from the first library and a fragment of about 600 bp was obtained from the third library. The composite sequence of 1154 nucleotides containing the complete predicted open reading frame of DRR-1 is shown in Figure 1. The open reading frame codes for a 337 amino acid protein (Figure 2) with a predicted molecular mass of 38.7 kD. The protein sequence contains all the characteristic
25 features of G protein-coupled receptors: seven hydrophobic helices likely to represent transmembrane domains, potential glycosylation site at the N-terminal extracellular domain (position 30) and a conserved NPXXY sequence at position 285-289.

Example 2: Cloning of Human DRR Receptor Genes

Two approaches were used to identify and clone novel human DNA sequences homologous and/or related to the rat DRR-1 gene. First, a human genomic library was screened in the lambda vector, Fix II, (Stratagene Cat.# 946203). Approximately 106 human genomic clones were plated and transferred onto nitrocellulose membranes for hybridization with the full length, 32P labeled, rat DRR-1 sequence as a probe. The hybridization was performed at 42 °C, overnight. The filters were washed at room temperature at low stringency (1X SSC/ 0.1% SDS) to allow detection of related but not necessarily identical sequences.

The inserted human DNA present in positive phages was amplified by PCR using the "Expand PCR kit" from Boehringer-Mannheim under conditions allowing accurate amplification of very large fragments of DNA. These long fragments of DNA were digested with various restriction enzymes and subcloned into a plasmid vector. The portions of these clones which hybridized with the rat DRR-1 gene probe were sequenced using the ABI cycle sequencing kit.

A second approach to identifying novel human sequences related to DRR-1 involved the use of the polymerase chain reaction (PCR), performed on total human genomic DNA. Primers were synthesized based upon the human genomic clones described above and were as follows:

HML.H, 5'-GCAAGCTTTCTGAGCATGGATCCAACCGTC, SEQ ID 21 and
HML.Bg, 5'-CCCTCAGATCTCCAATTGCTTCCCGACAG, SEQ ID NO:22.

Amplification resulted in a fragments of approximately 1 kilobase containing the entire coding sequence of the human genes. These fragments obtained were subcloned into the pGEM-T (Promega) vector for DNA sequencing analysis.

Using the above strategies, six human clones were isolated:

clone 7, SEQ ID numbers 3 and 4;

clone 18, SEQ ID numbers 5 and 6;

clone 23, SEQ ID numbers 7 and 8;

5 clone 24, SEQ ID numbers 9 and 10;

clone 36, SEQ ID numbers 11 and 12; and

clone 40, SEQ ID numbers 13 and 14.

None of these clones contain introns and their alignment may be seen in Figure 3.

10

At the amino acid sequence level, the rat DRR-1 clone is 47% to 49% identical to the human clones.

At the nucleic acid level, the rat DRR-1 clone is 56% to 58% identical to the human
15 clones. The level of sequence identity within the human clones (7, 18, 23, 24, 36, 40) is very high, between 77% and 98% at the amino acid sequence level. All the human sequences were used as queries to search for homologies in public databases (Genbank, Swissprot, EST). No identical sequences were detected. The closest matches were to members of the mas oncogene family of proteins. The overall amino acid sequence
20 homology between rat DRR-1 and any of the isolated human genes varied from 47 to 50 %. However some stretches display a much higher level of sequence homology, particularly the regions encoding the putative transmembrane domain III and VII (TM3 and TM7) and the intracellular loops 2 and 3 where the homology between the rat sequence and its human homologue is around 80%.

25

Example 3: In Situ Hybridization Experiments

Preparation of Tissue: Adult male Sprague-Dawley rats (~300 g; Charles River, St-Constant, Quebec) were sacrificed by decapitation. Brain and spinal cord with dorsal root ganglia attached were removed, snap-frozen in isopentane at -40°C for 20 s and stored
30 at -80 °C. Frozen human brain, spinal cord and dorsal root ganglia were obtained from the

Brain and Tissue Bank for Developmental Disorders, University of Maryland at Baltimore, according to the strictest ethical guidelines. Frozen tissue was sectioned at 14 μ m in a Microm HM 500 M cryostat (Germany) and thaw-mounted onto ProbeOn Plus slides (Fisher Scientific, Montreal, Quebec). Sections were stored at -80°C prior to in situ hybridization.

Synthesis of Riboprobes: The plasmid pGemT-3b32 GPCR was linearized using either SacII and Not I restriction enzymes. Sense and antisense DRR riboprobes were transcribed in vitro using either T7 or SP6 RNA polymerases (Pharmacia Biotech), respectively in the presence of [35S]UTP (~800 Ci/mmol; Amersham, Oakville, Ontario). The plasmid pGemT-Clone 36 GPCR was linearized using SacII and Pst I restriction enzymes. Sense and antisense Clon36 riboprobes were transcribed in vitro using either SP6 or T7 RNA polymerases (Pharmacia Biotech), respectively in the presence of [35S]UTP. Following transcription, the DNA template was digested with DNase I (Pharmacia). Riboprobes were purified by phenol/chloroform/isoamyl alcohol extraction and precipitated in 70% ethanol containing ammonium acetate and tRNA. Quality of labeled riboprobes was verified by polyacrylamide-urea gel electrophoresis.

In situ Hybridization: Sections were postfixed in 4% paraformaldehyde (BDH, Poole, England) in 0.1 M phosphate buffer (pH 7.4) for 10 min at room temperature (RT) and rinsed in three changes of 2X standard sodium citrate buffer (SSC; 0.15 M NaCl, 0.015 M sodium citrate, pH 7.0). Sections were then equilibrated in 0.1 M triethanolamine, treated with 0.25% acetic anhydride in triethanolamine, rinsed in 2X SSC and dehydrated in an ethanol series (50-100%). Hybridization was performed in a buffer containing 75% formamide (Sigma, St-Louis, Mo), 600 mM NaCl, 10 mM Tris (pH 7.5), 1 mM EDTA, 1X Denhardt's solution (Sigma), 50 (g/ml denatured salmon sperm DNA (Sigma), 50 (g/ml yeast tRNA (Sigma), 10% dextran sulfate (Sigma), 20 mM dithiothreitol and [35S]UTP-labeled cRNA probes (10 X10⁶ cpm/ml) at 55°C for 18 h in humidified chambers. Following hybridization, slides were rinsed in 2X SSC at RT, treated with 20 (g/ml RNase IA (Pharmacia) in RNase buffer (10 mM Tris, 500 mM NaCl, 1 mM EDTA, pH 7.5) for 45

min at RT and washed to a final stringency of 0.1X SSC at 65 °C. Sections were then dehydrated and exposed to Kodak Biomax MR film for 21 days and/or dipped in Kodak NTB2 emulsion diluted 1:1 with distilled water and exposed for 4-6 weeks at 4°C prior to development and counterstaining with cresyl violet acetate (Sigma).

5

Results: Of all regions examined within the neuraxis of the rat, DRR-1 mRNA was exclusively expressed in dorsal root ganglia. High resolution emulsion autoradiography showed accumulations of silver grains exclusively over small and some medium size neurons. This unique and highly restricted distribution pattern for DRR-1 was confirmed in
10 the rat embryo. Sagittal section of an E17 rat fetus showed that DRR-1 mRNA is confined to DRGs. All other structures of the rat embryo were devoid of any specific hybridization signal reinforcing the highly selective nature of DRR-1 expression

The expression of human Clone 36 receptor was present in human fetal dorsal root
15 ganglia but not in spinal cord. Specific hybridization signal for Clone 36 was not detected in any of the human adult CNS tissues examined thus far. These include spinal cord, cortex, hippocampus, thalamus, substantia nigra and periaqueductal gray (data not shown). Presence of Clone 36 mRNA in adult DRGs remains to be examined. Standard controls in which additional spinal cord with DRG sections were hybridized with rat DRR-1 antisense
20 or Clone 36 sense 35S-labeled probes displayed no specific hybridization signal.

Example 4: Northern Blots

Commercial rat and human multiple Northern blots containing 2 g of polyA RNA from various tissues (Clontech) were used to determine the expression and distribution of the rat
25 DRR-1 message and its human homologues. Radioactively labeled probes were prepared as follows: twenty five ng of a 650 bp 3b-32 PCR fragment derived from rat DRR-1 (see Example 1) or human clone 36 were random-prime labeled using the Ready-to-Go DNA labeling kit (Pharmacia Biotech) and [32P]CTP (3000 Ci/mmol/Amersham). The blot was prehybridized for 1 hour at 68 °C using Expresshyb (Clontech) followed by hybridization
30 (2X10⁶ cpm/ml of probe) for one hour using the same conditions. Blots were washed at

room temperature in 2X SSC, 0.05% SDS for 30 min. followed by 3x washes in 0.2X SSC, 1 % SDS at 50 °C for 60 min. and exposed at -80 °C to Kodak Biomax film for 6 days.

- 5 **Expression and Distribution of rat DRR-1:** All the rat tissues studied (heart, brain, spleen, lung, skeletal muscle, kidney and testis) were negative for the expression of DRR-1 following 2 weeks exposure whereas rat genomic Southern analysis revealed a 1.1 kb band when probed with the same cDNA fragment.
- 10 **Expression and Distribution of Human Clone 36:** Northern blots containing RNA from various human tissues were probed with a radio-labeled DNA fragment from clone 36. All the human tissues studied (human fetal brain, lung, liver and kidney and adult human cerebellum, cerebral cortex, medulla, occipital pole, frontal lobe, temporal lobe, putamen, spinal cord, amygdala, caudate nucleus, corpus callosum, hippocampus, total brain,
- 15 subthalamic nucleus and thalamus) were negative for the expression of this receptor following 2 weeks exposure.

Example 5: Calcium Signaling in Response to Angiotensin I-III

- The coding sequence of human clone 24 was transferred into a pcDNA3 vector and
- 20 modified to add a haemagglutinin tag at the C-terminus of the receptor sequence. This clone, designated as pcDNA3-HML-HA24 was transfected into HEK293 cells using a modified CaCl_2 method (Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press (1989)). The cells were maintained in culture medium at 37 °C, 5% CO_2 and diluted 10 fold every 3 days.

25

- The cells were inoculated in 90 mm tissue culture dishes (5 x 10⁵ cells per flask) in Dulbecco's Modified Essential Medium (DMEM, Gibco BRL), supplemented with 10% fetal bovine serum (FBS), 100 U/ml penicillin, 100 µg/ml streptomycin and 0.25 µg/ml fungizone. One day after inoculation, cells were transiently transfected with 30 µg of
- 30 plasmid DNA per dish. The cells were harvested 48 hours post transfection for analysis.

The expression of the gene was first checked by immunoprecipitation and western blotting with an anti-haemagglutinin antibody. A protein of approximately 43 KD was detected in stably as well as transiently transfected HEK293 cells, but not in control cells.

- 5 Stably transtected HEK293 cells were obtained after approximately 21 days of selection in culture medium containing 800 µg/ml G418. Calcium signaling measurement was performed with one of these stably transfected cell line, 293/pcDNA3-HML-HA24. The cells were grown on a 24 mm round glass cover slides to 50-70% confluence. After rinsing the cells with 1.8 NBS buffer (135 mM NaCl, 5 mM KCl, 1.2 mM MgCl₂, 1.8 mM
- 10 CaCl₂, 5 mM glucose and 10 mM HEPES, pH 7.3), the cells were incubated for one hour at room temperature in the presence of 0.5 ml of 3.5 µM FURA-2 AM (Molecular Probe, F-1221) diluted in 1.8 NBS. The cells were then rinsed three times with 1.8 NBS and incubated for a further 30 minutes at room temperature. The calcium displacement was measured using a PTI (Photon Technology International) D104 photometer equipped with a
- 15 PTI Delta RAM High speed multiwavelength illuminator, a PTI SC500 Shutter controller, a PTI LPS220 ARC lamp supply and the PTI FELIX software, v.1.2. Groups of 2 to 8 cells were chosen and isolated with the photometer diaphragm. The cells were exposed to 340 and 380 nm light and the 510 nm light emitted by the cells was recorded. Angiotensin I, II and III, were added successively - in various order from one experiment to the next -
- 20 followed by bradykinin as a positive control. Upon stimulation with angiotensin II and angiotensin III, a significant response was obtained. Addition of angiotensin I produced no response.

- All references cited herein are fully incorporated by reference. Having now fully
- 25 described the invention, it will be understood by one of skill in the art that the invention may be performed within a wide and equivalent range of conditions, parameters and the like, without affecting the spirit or scope of the invention or any embodiment thereof.

Applicant's or agent's file reference	N 1807-1 WO	International application No.
--	-------------	-------------------------------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>6</u> . line <u>8-9</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ)	
Address of depositary institution (including postal code and country) Mascheroder Weg 1b D-38124 Braunschweig Germany	
Date of deposit 27 November 1997	Accession Number DSM 11877
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
In respect of all designated states in which such action is possible and to the extent that it is legally permissible under the law of the designated state, it is requested that a sample of the deposited micro-organism be made available only by the issue thereof to an independent expert, in accordance with the relevant patent legislation, e.g. Rule 28(4) EPC, and generally similar provisions <i>mutatis mutandis</i> for any other designated state.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

SUBSTITUTE SHEET (RULE 26)

CLAIMS

1. A protein, except as existing in nature, comprising the amino acid sequence consisting functionally of rat dorsal root receptor 1 (DRR-1) as shown in SEQ ID NO:1.
5
2. A substantially pure protein according to claim 1, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:1.
3. A substantially pure polynucleotide encoding a protein comprising the amino acid
10 sequence consisting functionally of rat DRR-1 as shown in SEQ ID NO:1.
4. The polynucleotide of claim 3, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:1.
- 15 5. The polynucleotide of claim 4, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:2.
6. A protein, except as existing in nature, comprising the amino acid sequence consisting functionally of human dorsal root receptor 1 (DRR-1) as shown in SEQ ID NO:3.
20
7. A substantially pure protein according to claim 6, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:3.
8. A substantially pure polynucleotide encoding a protein comprising the amino acid
25 sequence consisting functionally of human DRR-1 as shown in SEQ ID NO:3.
9. The polynucleotide of claim 8, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:3.

10. The polynucleotide of claim 9, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:4.
11. A protein, except as existing in nature, comprising the amino acid sequence consisting functionally of human dorsal root receptor 2 (DRR-2) as shown in SEQ ID NO:5.
12. A substantially pure protein according to claim 11, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:5.
13. A substantially pure polynucleotide encoding a protein comprising the amino acid sequence consisting functionally of human DRR-2 as shown in SEQ ID NO:5.
14. The polynucleotide of claim 13, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:5.
15. The polynucleotide of claim 14, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:6.
16. A protein, except as existing in nature, comprising the amino acid sequence consisting functionally of human dorsal root receptor 3 (DRR-3) as shown in SEQ ID NO:7.
17. A substantially pure protein according to claim 16, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:7.
18. A substantially pure polynucleotide encoding a protein comprising the amino acid sequence consisting functionally of human DRR-3 as shown in SEQ ID NO:7.
19. The polynucleotide of claim 18, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:7.

20. The polynucleotide of claim 19, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:8.
21. A protein, except as existing in nature, comprising the amino acid sequence consisting
5 functionally of human dorsal root receptor 4 (DRR-4) as shown in SEQ ID NO:9.
22. A substantially pure protein according to claim 21, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:9.
- 10 23. A substantially pure polynucleotide encoding a protein comprising the amino acid sequence consisting functionally of human DRR-4 as shown in SEQ ID NO:9.
24. The polynucleotide of claim 23, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:9.
- 15 25. The polynucleotide of claim 24, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:10.
26. A protein, except as existing in nature, comprising the amino acid sequence consisting
20 functionally of human dorsal root receptor 5 (DRR-5) as shown in SEQ ID NO:11.
27. A substantially pure protein according to claim 26, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:11.
- 25 28. A substantially pure polynucleotide encoding a protein comprising the amino acid sequence consisting functionally of human DRR-5 as shown in SEQ ID NO:11.
29. The polynucleotide of claim 28, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:11.
- 30

30. The polynucleotide of claim 29, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:12.
31. A protein, except as existing in nature, comprising the amino acid sequence consisting
5 functionally of human dorsal root receptor 6 (DRR-6) as shown in SEQ ID NO:13.
32. A substantially pure protein according to claim 31, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:13.
- 10 33. A substantially pure polynucleotide encoding a protein comprising the amino acid sequence consisting functionally of human DRR-6 as shown in SEQ ID NO:13.
34. The polynucleotide of claim 33, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:13.
- 15 35. The polynucleotide of claim 34, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:14.
36. An antibody made by a process comprising the step of injecting a pharmaceutically
20 acceptable preparation comprising the protein of anyone of claims 1, 2, 6, 7, 11, 12, 16, 17, 21, 22, 26, 27, 31 or 32, into an animal capable of producing said antibody.
37. An antibody that binds specifically to anyone of the proteins of claims 1, 2, 6, 7, 11, 12, 16, 17, 21, 22, 26, 27, 31 or 32
- 25 38. A vector for expressing rat DRR-1, comprising the polynucleotide of either one of claim 3 or 4.

39. A vector for expressing anyone of
- (i) human DRR-1, comprising a polynucleotide of claim 8 or 9;
 - (ii) human DRR-2, comprising a polynucleotide of claim 13 or 14;
 - 5 (iii) human DRR-3, comprising a polynucleotide of claim 18 or 19;
 - (iv) human DRR-4, comprising a polynucleotide of claim 23 or 24;
 - (v) human DRR-5, comprising a polynucleotide of claim 28 or 29;
 - (vi) human DRR-6, comprising a polynucleotide of claim 33 or 34.
- 10 40. A host cell transformed with a vector according to claim 38 or 39.
41. Recombinant rat DRR-1, human DRR-1, human DRR-2, human DRR-3,
human DRR-4, human DRR-5, human DRR-6, produced by the host cell of claim 40.
- 15 42. A method for assaying a test compound for its ability to bind or to activate a G
protein-coupled dorsal root ganglia specific receptor (DRR), comprising:
- a) incubating a source containing DRR but substantially free of other G protein-
coupled receptors, with
 - i) a ligand known to bind to DRR;
 - 20 ii) said test compound; and
 - b) determining the extent to which said ligand binding is displaced by said test
compound.

25

30

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT: Astra Pharma Inc. Canada

(ii) TITLE OF INVENTION: Novel receptor

10

(iii) NUMBER OF SEQUENCES: 22

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Astra AB, Patent Department

(B) STREET: S-151 85 Södertälje

15

(C) CITY: Södertälje

(D) STATE:

(E) COUNTRY: Sweden

(F) ZIP: none

20

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

30

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 46-8 553 26000

(B) TELEFAX: 46-8 553 28820

5

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 337 amino acids

10

(B) TYPE: amino acid

(C) STRANDEDNESS: not relevant

(D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

15

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Met Val Cys Val Leu Arg Asp Thr Thr Gly Arg Phe Val Ser Met Asp

1

5

10

15

25

Pro Thr Ile Ser Ser Leu Ser Thr Glu Ser Thr Thr Leu Asn Lys Thr

20

25

30

Gly His Pro Ser Cys Arg Pro Ile Leu Thr Leu Ser Phe Leu Val Pro

35

40

45

30

SUBSTITUTE SHEET (RULE 26)

Ile Ile Thr Leu Leu Gly Leu Ala Gly Asn Thr Ile Val Leu Trp Leu
 50 55 60

Leu Gly Phe Arg Met Arg Arg Lys Ala Ile Ser Val Tyr Val Leu Asn
 5 65 70 75 80

Leu Ser Leu Ala Asp Ser Phe Phe Leu Cys Cys His Phe Ile Asp Ser
 85 90 95

Leu Met Arg Ile Met Asn Phe Tyr Gly Ile Tyr Ala His Lys Leu Ser
 10 100 105 110

Lys Glu Ile Leu Gly Asn Val Ala Phe Ile Pro Tyr Ile Ser Gly Leu
 115 120 125

Ser Ile Leu Ser Ala Ile Ser Thr Glu Arg Cys Leu Ser Val Leu Trp
 130 135 140

Pro Ile Trp Tyr His Cys His Arg Pro Arg Asn Met Ser Ala Ile Ile
 145 150 155 160

Cys Val Leu Ile Trp Val Leu Ser Phe Leu Met Gly Ile Leu Asp Trp
 165 170 175

Phe Phe Ser Gly Phe Leu Gly Glu Thr His His His Leu Trp Lys Asn
 180 185 190

Val Asp Phe Ile Val Thr Ala Phe Leu Ile Phe Leu Phe Met Leu Leu
 195 200 205

30

SUBSTITUTE SHEET (RULE 26)

Phe Gly Ser Ser Leu Ala Leu Leu Val Arg Ile Leu Cys Gly Ser Arg
210 215 220

Arg Lys Pro Leu Ser Arg Leu Tyr Val Thr Ile Ser Leu Thr Val Met
5 225 230 235 240

Val Tyr Leu Ile Cys Gly Leu Pro Leu Gly Leu Tyr Leu Phe Leu Leu
245 250 255

Tyr Trp Phe Gly Ile His Leu His Tyr Pro Phe Cys His Ile Tyr Gln
10 260 265 270

Val Thr Val Leu Leu Ser Cys Val Asn Ser Ser Ala Asn Pro Ile Ile
275 280 285

Tyr Phe Leu Val Gly Ser Phe Arg His Arg Lys Lys His Arg Ser Leu
15 290 295 300

Lys Met Val Leu Lys Arg Ala Leu Glu Glu Thr Pro Glu Glu Asp Glu
20 305 310 315 320

Tyr Thr Asp Ser His Val Gln Lys Pro Thr Glu Ile Ser Glu Arg Arg
325 330 335

Cys
25

SUBSTITUTE SHEET (RULE 26)

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 1011 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ATGGTTTGTG TTCTCAGGGA CACTACTGGA AGATTGTGA GCATGGATCC AACCATCTCA 60
20 TCCCTCAGTA CAGAATCTAC AACACTGAAT AAAACTGGTC ATCCCAGTTG CAGGCCAATC 120
CTCACCTGT CCTTCCTGGT CCCCATCATC ACCCTGCTTG GATTGGCAGG AAACACCATT 180
GTACTCTGGC TCTTGGGATT CCGCATGCGC AGGAAAGCCA TCTCAGTCTA CGTCCTCAAC 240
25 CTGTCTCTGG CAGACTCCTT CTTCTCTGTC TGCCATTTTA TTGACTCTCT GATGCGGATC 300
ATGAACTTCT ATGGCATCTA TGCCCATAAA TTAAGCAAAG AAATCTTAGG CAATGTAGCA 360
30 TTCATTCCCT ATATCTCAGG CCTGAGCATC CTCAGTGCTA TCAGCACGGA GCGCTGCCTG 420

SUBSTITUTE SHEET (RULE 26)

5
10
15
20

TCTGTATTGT GGCCAATCTG GTACCACTGC CACCGCCCAA GAAACATGTC AGCTATTATA 480

TGTGTTCTAA TCTGGGTCTT GTCTTTCTC ATGGGCATCC TTGACTGGTT TTTCTCAGGA 540

TTCTCTGGGTG AGACTCACCA TCATTTGTGG AAAAATGTTG ACTTTATTGT AACTGCATTT 600

CTGATTTTTT TATTTATGCT TCTCTTTGGG TCCAGTCTGG CGCTACTGGT GAGGATCCTC 660

TGTGGTTCCA GACGGAAACC ACTGTCCAGG CTGTACGTTA CAATCTCTCT CACAGTGATG 720

GTCTACCTCA TCTGCGGCCT GCCTCTCGGG CTTTACTTGT TCCTGCTATA TTGGTTTGGG 780

ATCCATTTAC ATTATCCCTT TTGTCACATT TACCAAGTTA CTGTGCTCCT GTCCTGTGTG 840

AACAGCTCTG CCAACCCCAT CATTTACTTC CTTGTAGGGT CCTTTAGGCA CCGTAAAAAG 900

CATCGGTCCC TCAAAATGGT TCTTAAAAGG GCTCTGGAGG AGACTCCTGA GGAGGATGAA 960

TATACAGACA GCCATGTTCA GAAACCCACT GAGATCTCAG AAAGGAGATG T 1011

(2) INFORMATION FOR SEQ ID NO:3:

- 15 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 322 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

5 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met Asp Pro Thr Ile Pro Val Leu Gly Thr Lys Leu Thr Pro Ile Asn
10 1 5 10 15
Gly Arg Glu Glu Thr Pro Cys Tyr Asn Gln Thr Leu Ser Phe Thr Gly
20 25 30
Leu Thr Cys Ile Ile Ser Leu Val Ala Leu Thr Gly Asn Ala Val Val
15 35 40 45
Leu Trp Leu Leu Gly Cys Arg Met Arg Arg Asn Ala Val Ser Ile Tyr
50 55 60
Ile Leu Asn Leu Val Ala Ala Asn Phe Leu Phe Leu Ser Gly His Ile
20 65 70 75 80
Ile Phe Ser Pro Leu Pro Leu Ile Asn Ile Arg His Pro Ile Ser Lys
25 85 90 95
Ile Leu Ser Pro Val Met Thr Phe Pro Tyr Phe Ile Gly Leu Ser Met
100 105 110

SUBSTITUTE SHEET (RULE 26)

Leu Ser Ala Ile Ser Thr Glu Arg Cys Leu Ser Ile Leu Trp Pro Ile
 115 120 125

5 Trp Tyr His Cys Arg Arg Pro Arg Tyr Leu Ser Ser Val Met Cys Val
 130 135 140

Leu Leu Trp Ala Leu Ser Leu Leu Arg Ser Ile Leu Glu Trp Met Phe
 145 150 155 160

10 Cys Asp Phe Leu Phe Ser Gly Ala Asn Ser Val Trp Cys Glu Thr Ser
 165 170 175

Asp Phe Ile Thr Ile Ala Trp Leu Val Phe Leu Cys Val Val Leu Cys
 15 180 185 190

Gly Ser Ser Leu Val Leu Leu Val Arg Ile Leu Cys Gly Ser Arg Lys
 195 200 205

20 Met Pro Leu Thr Arg Leu Tyr Val Thr Ile Leu Leu Thr Val Leu Val
 210 215 220

Phe Leu Leu Cys Gly Leu Pro Phe Gly Ile Gln Trp Ala Leu Phe Ser
 225 230 235 240

25 Arg Ile His Leu Asp Trp Lys Val Leu Phe Cys His Val His Leu Val
 245 250 255

Ser Ile Phe Leu Ser Ala Leu Asn Ser Ser Ala Asn Pro Ile Ile Tyr
 30 260 265 270

SUBSTITUTE SHEET (RULE 26)

Phe Phe Val Gly Ser Phe Arg Gln Arg Gln Asn Arg Gln Asn Leu Lys

275

280

285

5 Leu Val Leu Gln Arg Ala Leu Gln Asp Thr Pro Glu Val Asp Glu Gly

290

295

300

Gly Gly Trp Leu Pro Gln Glu Thr Leu Glu Leu Ser Gly Ser Lys Leu

10

305

310

315

320

Glu Gln

15 (2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 969 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

30

SUBSTITUTE SHEET (RULE 26)

ATGGATCCAA CCATCCCACT CTTGGGTACA AAACGACAC CAATCAACGG ACGTGAGGAG 60

ACTCCTTGCT ACAACCAAAC CCGAGCTTC ACGGGGCTGA CGTGATCAT TTCCCTGTG 120

5 GCGCTGACAG GAAACGCGT TGTGCTCTGG CTCCTGGGCT GCCGCATGCG CAGGAACGCT 180

GTCTCCATCT ACATCCTCAA CCGGTGCGG GCCAACTTCC TCTTCCTTAG CGGCCACATT 240

ATATTTTCGC CGTTACCCCT CATCAATATC CGCCATCCCA TCTCCAAAAT CCTCAGTCCT 300

10 GTGATGACCT TTCCCTACTT TATAGGCCTA AGCATGCTGA GCGCCATCAG CACCGAGCGC 360

TGCCTGTCCA TCCTGTGGCC CATCTGGTAC CACTGCCGCC GCCCCAGATA CCTGTCATCG 420

15 GTCATGTGTG TCCTGCTCTG GGCCCTGTCC CTGCTGCCGA GTATCCTGGA GTGGATGTTT 480

TGTGACTTCC TGTTTAGTGG TGCTAATTCT GTTTGGTGTG AAACGTCAGA TTTCATTACA 540

ATCGCGTGGC TGGTTTTTTT ATGTGTGGTT CTCTGTGGGT CCAGCCTGGT CCTGCTGGTC 600

20 AGGATTCTCT GTGGATCCCG GAAGATGCCG CTGACCAGGC TGTACGTGAC CATCCTCCTC 660

ACAGTGCTGG TCTTCCTCCT CTGTGGCCTG CCCTTTGGCA TTCAGTGGGC CCTGTTTTCC 720

25 AGGATCCACC TGGATTGGAA AGTCTTATTT TGTCATGTGC ATCTAGTTTC CATTTTCCTG 780

TCCGCTCTTA ACAGCAGTGC CAACCCCATC ATTACTTCT TCGTGGGCTC CTTAGGCAG 840

CGTCAAAATA GGCAAAACCT GAAGCTGGTT CTCCAAAGGG CTCTGCAGGA CACGCCTGAG 900

30

GTGGATGAAG GTGGAGGGTG GCTTCCTCAG GAAACCCTGG AGCTGTCGGG AAGCAAATTG 960

GAGCAGTGA 969

5

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 322 amino acids

10

(B) TYPE: amino acid

(C) STRANDEDNESS: not relevant

(D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

15

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Asp Pro Thr Val Pro Val Leu Gly Thr Glu Leu Thr Pro Ile Asn

1

5

10

15

25

Gly Arg Glu Glu Thr Pro Cys Tyr Lys Gln Thr Leu Ser Phe Thr Gly

20

25

30

Leu Thr Cys Ile Val Ser Leu Val Ala Leu Thr Gly Asn Ala Val Val

35

40

45

30

SUBSTITUTE SHEET (RULE 26)

Leu Trp Leu Leu Gly Cys Arg Met Arg Arg Asn Ala Val Ser Ile Tyr
 50 55 60

5 Ile Leu Asn Leu Val Ala Ala Asp Phe Leu Phe Leu Ser Gly His Ile
 65 70 75 80

Ile Cys Ser Pro Leu Arg Leu Ile Asn Ile Ser His Pro Ile Ser Lys
 85 90 95

10 Ile Leu Ser Pro Val Met Thr Phe Pro Tyr Phe Ile Gly Leu Ser Met
 100 105 110

Leu Asn Ala Ile Ser Thr Glu Arg Cys Leu Ser Ile Leu Trp Pro Ile
 115 120 125

15 Trp Tyr His Cys Arg Arg Pro Arg Tyr Leu Ser Ser Val Met Cys Val
 130 135 140

Leu Leu Trp Ala Pro Ser Leu Leu Arg Ser Ile Leu Glu Trp Met Phe
 20 145 150 155 160

Cys Asp Phe Leu Phe Ser Gly Ala Asp Ser Val Arg Cys Glu Thr Ser
 165 170 175

25 Asp Phe Ile Thr Ile Ala Trp Leu Val Phe Leu Arg Val Val Leu Cys
 180 185 190

Gly Ser Ser Leu Val Leu Leu Val Arg Ile Leu Cys Gly Ser Arg Lys
 195 200 205

30

Met Pro Leu Thr Arg Leu Tyr Val Thr Ile Leu Leu Thr Val Leu Val

210

215

220

Phe Leu Leu Cys Gly Leu Pro Phe Gly Ile Gln Trp Ala Leu Phe Ser

5

225

230

235

240

Arg Ile His Leu Asp Trp Lys Val Leu Phe Cys His Val His Leu Val

245

250

255

10

Ser Ile Phe Leu Ser Ala Leu Asn Ser Ser Ala Asn Pro Ile Ile Tyr

260

265

270

Phe Phe Met Gly Ser Phe Arg Gln Leu Gln Asn Arg Lys Thr Leu Lys

15

275

280

285

Leu Val Leu Gln Arg Asp Leu Gln Asp Thr Pro Glu Val Asp Glu Gly

290

295

300

20

Gly Trp Trp Leu Pro Gln Glu Thr Leu Glu Leu Ser Gly Ser Lys Leu

305

310

315

320

Glu Ile

25

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 969 base pairs

(B) TYPE: nucleic acid

30

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 98/02348

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07K 14/72

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9504073 A1 (THE BOARD OF TRUSTEES OF THE LELAND STANFORD JUNIOR UNIVERSITY), 9 February 1995 (09.02.95)	1-35, 38-42
A	--	36-37
X	WO 9405695 A1 (NEW YORK UNIVERSITY), 17 March 1994 (17.03.94), See page 14, line 11 seq 52, claims	1-42
A	Dialog Information Service, file 154, Medline, Dialog accession no. 08044093, Medline accession no. 95047685, Brown NJ et al: "Gastrointestinal adaptation to enhanced small intestinal lipid exposure", Gut (ENGLAND) Oct 1994, 35 (10) p 1409-12	1-42
	--	

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

6 April 1999

Date of mailing of the international search report

14.04.99

Name and mailing address of the ISA/

Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Patrick Andersson
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

Information on patent family members

02/03/99

International application No.

PCT/SE 98/02348

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9504073	A1	09/02/95	US	5556780 A	17/09/96
WO	9405695	A1	17/03/94	AU	4855393 A	29/03/94
				US	5508384 A	16/04/96

Form PCT/ISA/210 (patent family annex) (July 1992)





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/72	A1	(11) International Publication Number: WO 99/32519 (43) International Publication Date: 1 July 1999 (01.07.99)
<p>(21) International Application Number: PCT/SE98/02348</p> <p>(22) International Filing Date: 16 December 1998 (16.12.98)</p> <p>(30) Priority Data: 9704836-7 22 December 1997 (22.12.97) SE</p> <p>(71) Applicant (for all designated States except MG US): ASTRA PHARMA INC. [CA/CA]; 1004 Middlegate Road, Mississauga, Ontario L4Y 1M4 (CA).</p> <p>(71) Applicant (for MG only): ASTRA AKTIEBOLAG [SE/SE]; S-151 85 Södertälje (SE).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): AHMAD, Sultan [IN/CA]; Astra Research Centre Montreal, 7171 Fredrick-Banting, St. Laurent, Quebec H4S 1Z9 (CA). BANVILLE, Denis [CA/CA]; 595 Lajeunesse, Ste-Dorothee, Quebec H7X 3K4 (CA). FORTIN, Yves [CA/CA]; 2985 Avenue Douglas, Montreal, Quebec H3R 2E2 (CA). LEMBO, Paola [CA/CA]; Astra Research Centre Montreal, 7171 Fredrick-Banting, St. Laurent, Quebec H4S 1Z9 (CA). O'DONELL, Dajan [CA/CA]; Astra Research Centre Montreal, 7171 Fredrick-Banting, St. Laurent, Quebec H4S</p>	<p>1Z9 (CA). SHEN, Shi-Hsiang [CA/CA]; 161 Charwell Crescent, Beaconsfield, Quebec H9W 1C2 (CA).</p> <p>(74) Agent: ASTRA AKTIEBOLAG; Intellectual Property, S-151 85 Södertälje (SE).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p>	
<p>(54) Title: NOVEL G PROTEIN-COUPLED RECEPTOR</p> <p>(57) Abstract</p> <p>The present invention is directed to novel G protein-coupled receptors that are found predominantly in the dorsal root ganglia. The invention encompasses both receptor proteins as well as nucleic acids encoding the proteins. Angiotension I and III effects Calcium signalling in Cells transformed with DNA encoding the receptor. In addition, the present invention is directed to methods and compositions which utilize the receptors.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

NOVEL G PROTEIN-COUPLED RECEPTOR

Field of the Invention

The present invention is in the general field of biological receptors and the various uses
5 that can be made of such receptors. More specifically, the invention relates to nucleic acids
encoding novel G protein-coupled receptors and to the receptors per se.

Background and Prior Art

G protein-coupled receptors (GPCRs) constitute a family of proteins sharing a common
10 structural organization characterized by an extracellular N-terminal end, seven hydrophobic
alpha helices putatively constituting transmembrane domains and an intracellular C-
terminal domain. GPCRs bind a wide variety of ligands that trigger intracellular signals
through the activation of transducing G proteins (*Caron, et al., Rec. Prog. Horm. Res.*
48:277-290 (1993); *Freedman et al., Rec. Prog. Horm. Res.* 51:319-353 (1996)).

15 More than 300 GPCRs have been cloned thus far and it is generally assumed that there
exist well over 1000 such receptors. Mechanistically, approximately 50-60% of all
clinically relevant drugs act by modulating the functions of various GPCRs (*Cudermann, et*
al., J. Mol. Med. 73:51-63 (1995)). Of particular interest are receptors located in dorsal root
20 ganglia. This region of the central nervous system is densely innervated with primary or
afferent sensory neurons involved in the transmission, modulation and sensation of pain.
Thus, receptors from this region may be used in assays for the identification of new agents
for anesthesia and analgesia

Summary of the Invention

25 The present invention is based upon the discovery of a novel G protein-coupled receptor
which is distinct from previously reported receptors in terms of structure and in being
expressed preferentially in dorsal root ganglia. One dorsal root receptor (DRR) has been
isolated and sequenced from the rat and six from the human. The rat receptor was given the

designation rDRR-1 and its amino acid sequence is shown as SEQ ID NO:1. The human receptors were designated as

hDRR-1 (SEQ ID NO:3);

hDRR-2 (SEQ ID NO:5);

5 hDRR-3 (SEQ ID NO:7):

hDRR-4 (SEQ ID NO:9);

hDRR-5 (SEQ ID NO:11); and

hDRR-6 (SEQ ID NO:13).

Unless otherwise specified, the term "DRR" as used herein refers to all of the receptors
10 from both human and rat .

In its first aspect, the invention is directed to proteins, except as existing in nature, comprising the amino acid sequence consisting functionally of a rat or human DRR as shown in SEQ ID NO:1, 3, 5, 7, 9, 11, or 13. The term "consisting functionally of" is
15 intended to include any receptor protein whose sequence has undergone additions, deletions or substitutions which do not substantially alter the functional characteristics of the receptor. Thus, the invention encompasses proteins having exactly the same amino acid sequence as shown in the sequence listing, as well as proteins with differences that are not substantial as evidenced by their retaining the basic, qualitative binding properties of
20 the DRR receptor. The invention further encompasses substantially pure proteins consisting essentially of a DRR amino acid sequence, antibodies that bind specifically to a DRR (i.e. that have at least a 100 fold greater affinity for the DRR than any other naturally occurring non-DRR protein), and antibodies made by a process involving the injection of pharmaceutically acceptable preparations of such proteins into an animal capable of
25 antibody production. In a preferred embodiment, monoclonal antibody to human or rat DRR is produced by injecting a pharmaceutically acceptable preparation of the receptor into a mouse and then fusing mouse spleen cells with myeloma cells.

The invention is also directed to a substantially pure polynucleotide encoding a protein
30 comprising the amino acid sequence consisting functionally of the sequence of rat DRR (as

shown in SEQ ID NO:1) or a human DRR (as shown in SEQ ID NOs 3, 5, 7, 9, 11 or 13). This aspect of the invention encompasses polynucleotides encoding proteins consisting essentially of the amino acid sequences shown in the sequence listing, expression vectors comprising such polynucleotides, and host cells transformed with such vectors. Also
5 included are the recombinant rat and human DRR proteins produced by host cells made in this manner.

Preferably, the polynucleotide encoding rat DRR has the nucleotide sequence shown in SEQ ID NO:2 and the polynucleotide encoding a human DRR has the nucleotide sequence
10 shown in SEQ ID NO: 3, 5, 7, 9, 11 or 13. It is also preferred that the vectors and host cells used for the expression of DRR contain these particular polynucleotides.

In another aspect, the present invention is directed to a method for assaying a test compound for its ability to bind to a rat or human DRR. The method is performed by
15 incubating a source of DRR with a ligand known to bind to the receptor and with the test compound. The source of the DRR should be substantially free of other types of G protein-coupled receptors, i.e. greater than 85% of such receptors present should correspond to the DRR. Upon completion of incubation, the ability of the test compound to bind to the DRR is determined by the extent to which ligand binding has been displaced. The rat DRR
20 should, preferably correspond to rDRR-1 as shown in SEQ ID NO:1. The human receptor should preferably be hDRR-1 (SEQ ID NO:3); hDRR-2 (SEQ ID NO:5); hDRR-3 (SEQ ID NO:7); hDRR-4 (SEQ ID NO:9); hDRR-5 (SEQ ID NO:11); or hDRR-6 (SEQ ID NO:13). Either transformed cells expressing recombinant DRR may be used in the assays or membranes can be prepared from the cells and used. Although not essential, the assay can
25 be accompanied by the determination of the activation of a second messenger pathway such as the adenyl cyclase pathway. This should help to determine whether a compound that binds to DRR is acting as an agonist or antagonist.

An alternative method for determining if a test compound is an agonist of any of the
30 DRRs disclosed herein is to use a cell signaling assay, e.g., an assay measuring either

intracellular adenylyl cyclase activity or intracellular calcium concentration. The test compound is incubated with cells expressing the DRR but substantially free of other G protein-coupled receptors, typically a cell transfected with an expression vector encoding the DRR. Test compounds that are agonists are identified by their causing a statistically significant change in the results obtained from the cell signaling assay when compared to control transfectants not exposed to test compound. For example, the cells exposed to the test compound may show a significant increase in adenylyl cyclase activity or in intracellular calcium concentration.

- 10 The invention also encompasses a method for determining if a test compound is an antagonist of a DRR which relies upon the known activation of G protein-coupled receptors that occurs when such receptors are expressed in large amounts. This method requires that DNA encoding the receptor be incorporated into an expression vector so that it is operably linked to a promoter and that the vector then be used to transfect an appropriate host. In order to produce sufficient receptor to result in constitutive receptor activation (i.e., activation in the absence of natural ligand), expression systems capable of copious protein production are preferred, e.g., the DRR DNA may be operably linked to a CMV promoter and expressed in COS or HEK293 cells. After transfection, cells with activated receptors are selected based upon their showing increased activity in a cell signaling assay relative to comparable cells that have either not been transfected or that have been transfected with a vector that is incapable of expressing functional DRR. Typically, cells will be selected either because they show a statistically significant increase in intracellular adenylyl cyclase activity or a statistically significant increase in intracellular calcium concentration. The selected cells are contacted with the test compound and the cell signaling assay is repeated to determine if this results in a decrease in activity relative to control cells not contacted with the test compound. For example, a statistically significant decrease in either adenylyl cyclase activity or calcium concentration relative to control cells would indicate that the test compound is an antagonist of the DRR. Any of the DRRs disclosed herein may be used in these assays.

- Assays for compounds interacting with a DRR may be performed by incubating a source containing the DRR but substantially free of other G protein-coupled receptors (e.g. a stably transformed cell) with angiotensin II or III in both the presence and absence of test compound and measuring the modulation of intracellular calcium concentration. A
- 5 significant increase or decrease in angiotensin-stimulated calcium displacement in response to test compound is indicative of an interaction occurring at the DRR. The receptors that may be used in these assays include rat DRR-1 and human DRR-1, DRR-2, DRR-3, DRR-4, DRR-5 and DRR-6.
- 10 In another aspect, the present invention is directed to a method for assaying a test compound for its ability to alter the expression of a rat or human DRR. This method is performed by growing cells expressing the DRR, but substantially free of other G protein-coupled receptors, in the presence of the test compound. Cells are then collected and the expression of the DRR is compared with expression in control cells grown under
- 15 essentially identical conditions but in the absence of the test compound. The rat receptor is preferably rDRR-1 and the human receptor may be DRR-1; DRR-2; DRR-3; DRR-4; DRR-5; or DRR-6.
- A preferred test compound is an oligonucleotide at least 15 nucleotides in length
- 20 comprising a sequence complimentary to the sequence of the DRR used in the assay.

Brief Description of the Drawings

Figure 1. Nucleotide sequence of rDDR-1: Clone 3B-32, encoding rDDR-1, was isolated from a rat genomic library using the Promoter Finder Walking Kit (see Methods, Clontech).

5

The cloned gene was deposited with the international depositary authority Deutsche Sammlung Von Mikroorganismen Und Zellkulturen GmbH at the address Mascheroder Weg 1 B, D-3300 Braunschweig, Germany. The deposit was made on November 27, 1997 and was given the accession number DSM 11877.

10

Figure 2. Deduced amino acid sequence of DDR-1: Clone 3B-32 codes for a 337 amino acid protein. The amino acid sequence begins with the first ATG in the nucleotide sequence.

15 Figure 3. Alignment of the deduced amino acid sequences of clone 3B-32 (rDDR-1) with its five most homologous sequences. The boxed and shaded residues are the ones that are identical to the rDDR-1 sequence.

Figure 4. Amino acid alignment of the human DDR homologs: The amino acid sequence of all 6 human homologs of rDDR-1 (hDDR-1; hDDR-2; hDDR-3; hDDR-4; hDDR-5; and hDDR-6) are aligned. The amino acid residues differing from the clone 36 (HUMAN36.PR) are boxed. The degree of identity among these sequences ranges from 77% to almost 100%.

25 Definitions

The description that follows uses a number of terms that refer to recombinant DNA technology. In order to provide a clear and consistent understanding of the specification and claims, including the scope to be given such terms, the following definitions are provided.

30

Cloning vector: A plasmid or phage DNA or other DNA sequence which is able to replicate autonomously in a host cell, and which is characterized by one or a small number of restriction endonuclease recognition sites. A foreign DNA fragment may be spliced into the vector at these sites in order to bring about the replication and cloning of the fragment.

5 The vector may contain a marker suitable for use in the identification of transformed cells. For example, markers may provide tetracycline resistance or ampicillin resistance.

Expression vector: A vector similar to a cloning vector but which is capable of inducing the expression of the DNA that has been cloned into it, after transformation into a host.

10 The cloned DNA is usually placed under the control of (i.e., operably linked to) certain regulatory sequences such as promoters or enhancers. Promoter sequences may be constitutive, inducible or repressible.

Substantially pure: As used herein, "substantially pure" means that the desired product is essentially free from contaminating cellular components. A "substantially pure" protein or nucleic acid will typically comprise at least 85% of a sample, with greater percentages being preferred. Contaminants may include proteins, carbohydrates or lipids. One method for determining the purity of a protein or nucleic acid is by electrophoresing a preparation in a matrix such as polyacrylamide or agarose. Purity is evidenced by the appearance of a

15 single band after staining. Other methods for assessing purity include chromatography and analytical centrifugation.

20

Host: Any prokaryotic or eukaryotic cell that is the recipient of a replicable expression vector or cloning vector is the "host" for that vector. The term encompasses prokaryotic or eukaryotic cells that have been engineered to incorporate a desired gene on its chromosome or in its genome. Examples of cells that can serve as hosts are well known in the art, as are techniques for cellular transformation (see e.g. Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd ed. Cold Spring Harbor (1989)).

25

30 **Promoter:** A DNA sequence typically found in the 5' region of a gene, located proximal

to the start codon. Transcription is initiated at the promoter. If the promoter is of the inducible type, then the rate of transcription increases in response to an inducing agent.

Complementary Nucleotide Sequence: A complementary nucleotide sequence, as used
5 herein, refers to the sequence that would arise by normal base pairing. For example, the nucleotide sequence 5'-AGAC-3' would have the complementary sequence 5'-GTCT-3'.

Expression: Expression is the process by which a polypeptide is produced from DNA. The process involves the transcription of the gene into mRNA and the translation of this
10 mRNA into a polypeptide.

Detailed Description of the Invention

The present invention is directed to DRR receptor proteins, genetic sequences coding for
15 the receptors, a method for assaying compounds for binding to DRR receptors and a method for assaying compounds for their ability to alter DRR expression. The receptors and their nucleic acids are defined by their structures (as shown in figures 1, 2 and 4; and SEQ ID numbers 1-14).

20 It will be understood that the present invention encompasses not only sequences identical to those shown in the figures and sequence listing, but also sequences that are essentially the same and sequences that are otherwise substantially the same and which result in a receptor retaining the basic binding characteristics of the DRR. For example, it is well known that techniques such as site-directed mutagenesis may be used to introduce
25 variations in a protein's structure. Variations in a DRR protein introduced by this or some similar method are encompassed by the invention provided that the resulting receptor retains the basic qualitative binding characteristics of the unaltered DRR. Thus, the invention relates to proteins comprising amino acid sequences consisting functionally of the sequence of SEQ ID NO:1 (rat) and SEQ ID numbers 3, 5, 7, 9, 11 and 14 (human).

I. Nucleic Acid Sequences Coding for DRR

DNA sequences coding for DRRs are expressed exclusively, or at least highly preferentially, in dorsal root ganglia and these ganglia may serve as a source for the isolation of nucleic acids coding for the receptors. In addition, cells and cell lines that
5 express a rat or human DRR may serve as a source for nucleic acid. These may either be cultured cells that have not undergone transformation or cell lines specifically engineered to express recombinant DRR.

In all cases, poly A+ mRNA is isolated from the dorsal root ganglia, reverse transcribed
10 and cloned. The cDNA library thus formed may then be screened using probes derived from the sequences shown in the accompanying sequence listing as SEQ ID number 2, 4, 6, 8, 10, 12 or 14, depending upon the particular DRR being isolated. Probes should typically be at least 14 bases in length and should be derived from a portion of the DRR sequence that is poorly conserved (see Figures 3 and 4). Screening can also be performed using
15 genomic libraries with one DRR gene, or a portion of the gene, serving as a probe in the isolation of other DRR genes. For example, full length rDRR-1 may be labeled and used to screen a human genomic library for the isolation of hDRR-1, hDRR-2 etc. (see Examples section).

20 Alternatively genomic DNA libraries can be used to isolate DRR genes by performing PCR amplifications with primers located at either end of genes (see Examples section for a description of procedures). For example, human genomic DNA may be amplified using the primers:

25 5'-GCAAGCTTTCTGAGCATGGATCCAACCGTC, and
5'-CCCTCAGATCTCCAATTTGCTTCCCGACAG.

This will serve to amplify all six of the human DRR genes identified herein as hDRR-1; hDRR-2; hDRR-3; hDRR-4; hDRR-5; and hDRR-6. These may then be cloned into an appropriate vector, e.g. pGEM-T (Promega), for DNA sequence analysis.

5 II. Antibodies to Rat and Human DRRs

The present invention is also directed to antibodies that bind specifically to a rat or human DRR and to a process for producing such antibodies. Antibodies that "bind specifically to a DRR" are defined as those that have at least a one hundred fold greater affinity for the DRR than for any other protein. The process for producing such antibodies may involve
10 either injecting the DRR protein itself into an appropriate animal or, preferably, injecting short peptides made to correspond to different regions of the DRR. The peptides should be at least five amino acids in length and should be selected from regions believed to be unique to the particular DRR protein being targeted. Thus, highly conserved transmembrane regions should generally be avoided in selecting peptides for the generation
15 of antibodies. Methods for making and detecting antibodies are well known to those of skill in the art as evidenced by standard reference works such as: (*Harlow et al., Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory, N.Y. (1988)*); (*Klein, Immunology: The Science of Self-Nonself Discrimination (1982)*); (*Kennett, et al., Monoclonal Antibodies and Hybridomas: A New Dimension in Biological Analyses (1980)*);
20 and (*Campbell, "Monoclonal Antibody Technology," in Laboratory Techniques in Biochemistry and Molecular Biology, (1984)*).

"Antibody," as used herein, is meant to include intact molecules as well as fragments which retain their ability to bind to antigen (e.g., Fab and F(ab)₂ fragments). These fragments are
25 typically produced by proteolytically cleaving intact antibodies using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab)₂ fragments). The term "antibody" also refers to both monoclonal antibodies and polyclonal antibodies. Polyclonal antibodies are derived from the sera of animals immunized with the antigen. Monoclonal antibodies can be prepared using hybridoma technology (*Kohler, et al., Nature 256:495 (1975)*); (*Hammerling, et al., in: Monoclonal Antibodies and T-Cell Hybridomas. Elsevier,*
30

M.Y., pp. 563-681 (1981)). In general, this technology involves immunizing an animal, usually a mouse, with either intact DRR or a fragment derived from the DRR. The splenocytes of the immunized animals are extracted and fused with suitable myeloma cells, e.g., SP2O cells. After fusion, the resulting hybridoma cells are selectively maintained in
5 HAT medium and then cloned by limiting dilution (Wands, et al., *Gastroenterology* 80:225-232 (1981)). The cells obtained through such selection are then assayed to identify clones which secrete antibodies capable of binding to the DRR.

The antibodies, or fragments of antibodies, of the present invention may be used to detect
10 the presence of DRR protein using any of a variety of immunoassays. For example, the antibodies may be used in radioimmunoassays or in immunometric assays, also known as "two-site" or "sandwich" assays (see *Chard, T., "An Introduction to Radioimmune Assay and Related Techniques," in Laboratory Techniques in Biochemistry and Molecular Biology, North Holland Publishing Co., N.Y. (1978)*). In a typical immunometric assay, a
15 quantity of unlabeled antibody is bound to a solid support that is insoluble in the fluid being tested, e.g., blood, lymph, cellular extracts, etc. After the initial binding of antigen to immobilized antibody, a quantity of detectably labeled second antibody (which may or may not be the same as the first) is added to permit detection and/or quantitation of bound antigen (see e.g. *Radioimmune Assay Method, Kirkham et al., ed., pp. 199-206. E & S. Livingstone, Edinburgh (1970)*). Many variations of these types of assays are known in the
20 art and may be employed for the detection of the DRR.

Antibodies to a rat or human DRR may also be used in the purification of either the intact receptor or fragments of the receptor (see generally, *Dean et al., Affinity Chromatography, A Practical Approach, IRL Press (1986)*). Typically, antibody is immobilized on a
25 chromatographic matrix such as Sepharose 4B. The matrix is then packed into a column and the preparation containing the DRR desired is passed through under conditions that promote binding, e.g., under conditions of low salt. The column is then washed and bound DRR is eluted using a buffer that promotes dissociation from antibody, e.g., buffer having

an altered pH or salt concentration. The eluted DRR may be transferred into a buffer of choice, e.g., by dialysis, and either stored or used directly.

III. Radioligand Assay for Receptor Binding

5 One of the main uses for DRR nucleic acids and recombinant proteins is in assays designed to identify agents capable of binding to DRR receptors. Such agents may either be agonists, mimicking the normal effects of receptor binding, or antagonists, inhibiting the normal effects of receptor binding. Of particular interest is the identification of agents which bind to the DRR and modulate adenylyl cyclase activity in the cells. These agents have
10 potential therapeutic application as either analgesics or anesthetics.

In radioligand binding assays, a source of DRR is incubated together with a ligand known to bind to the receptor and with the compound being tested for binding activity. The preferred source for DRR is cells, preferably mammalian cells, transformed to recombinantly express the receptor. The cells selected should not express a substantial
15 amount of any other G protein-coupled receptors that might bind to ligand and distort results. This can easily be determined by performing binding assays on cells derived from the same tissue or cell line as those recombinantly expressing DRR but which have not undergone transformation.

20 The assay may be performed either with intact cells or with membranes prepared from the cells (*see e.g. Wang, et al., Proc. Natl. Acad. Sci. U.S.A. 90:10230-10234 (1993)*). The membranes are incubated with a ligand specific for the DRR receptor and with a preparation of the compound being tested. After binding is complete, receptor is separated from the solution containing ligand and test compound, e.g. by filtration, and the amount of
25 binding that has occurred is determined. Preferably, the ligand used is detectably labeled with a radioisotope such as ¹²⁵I. However, if desired, fluorescent or chemiluminescent labels can be used instead. Among the most commonly used fluorescent labeling compounds are fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde and fluorescamine. Useful chemiluminescent compounds
30 include luminol, isoluminol, theromatic acridinium ester, imidazole, acridinium salt, and

oxalate ester. Any of these agents which can be used to produce a ligand suitable for use in the assay.

Nonspecific binding may be determined by carrying out the binding reaction in the presence of a large excess of unlabeled ligand. For example, labeled ligand may be incubated with receptor and test compound in the presence of a thousandfold excess of unlabeled ligand. Nonspecific binding should be subtracted from total binding, i.e. binding in the absence of unlabeled ligand, to arrive at the specific binding for each sample tested. Other steps such as washing, stirring, shaking, filtering and the like may be included in the assays as necessary. Typically, wash steps are included after the separation of membrane-bound ligand from ligand remaining in solution and prior to quantitation of the amount of ligand bound, e.g., by counting radioactive isotope. The specific binding obtained in the presence of test compound is compared with that obtained in the presence of labeled ligand alone to determine the extent to which the test compound has displaced receptor binding.

In performing binding assays, care must be taken to avoid artifacts which may make it appear that a test compound is interacting with the DRR receptor when, in fact, binding is being inhibited by some other mechanism. For example, the compound being tested should be in a buffer which does not itself substantially inhibit the binding of ligand to DRR and should, preferably, be tested at several different concentrations. Preparations of test compound should also be examined for proteolytic activity and it is desirable that antiproteases be included in assays. Finally, it is highly desirable that compounds identified as displacing the binding of ligand to DRR receptor be reexamined in a concentration range sufficient to perform a Scatchard analysis on the results. This type of analysis is well known in the art and can be used for determining the affinity of a test compounds for receptor (see e.g., Ausubel, et al., *Current Protocols in Molecular Biology*, 11.2.1-11.2.19 (1993); *Laboratory Techniques and Biochemistry and Molecular Biology*, Work, et al., ed., N.Y. (1978) etc.). Computer programs may be used to help in the analysis of results (see e.g., Munson, P., *Methods Enzymol.* 92:543-577 (1983); McPherson, G.A., *Kinetic, EBDA*

Ligand, Lowry-A Collection of Radioligand Binding Analysis Programs, Elsevier-Biosoft, U.K. (1985)).

The activation of receptor by the binding of ligand may be monitored using a number of
5 different assays. For example, adenylyl cyclase assays may be performed by growing cells in
wells of a microtiter plate and then incubating the various wells in the presence or absence
of test compound. cAMP may then be extracted in ethanol, lyophilized and resuspended in
assay buffer. Assay of cAMP thus recovered may be carried out using any method for
determining cAMP concentration, e.g. the Biotrack cAMP Enzyme-immunoassay System
10 (Amersham) or the Cyclic AMP [3H] Assay System (Amersham). Typically, adenylyl
cyclase assays will be performed separately from binding assays, but it may also be
possible to perform binding and adenylyl cyclase assays on a single preparation of cells.
Other "cell signaling assays" that can be used to monitor receptor activity are described
below.

15

IV. Identification of DRR Agonists and Antagonists Using Cell Signaling Assays

DRRs may also be used to screen for drug candidates using cell signaling assays. To
identify DRR agonists, the DNA encoding a receptor is incorporated into an expression
vector and then transfected into an appropriate host. The transformed cells are then
20 contacted with a series of test compounds and the effect of each is monitored. Among the
assays that can be used are assays measuring cAMP production (see discussion above),
assays measuring the activation of reporter gene activity, or assays measuring the
modulation of the binding of GTP-gamma-S.

25 Cell signaling assays may also be used to identify DRR antagonists. G protein-coupled
receptors can be put in their active state even in the absence of their cognate ligand by
expressing them at very high concentration in a heterologous system. For example, receptor
may be overexpressed using the baculovirus infection of insect Sf9 cells or a DRR gene
may be operably linked to a CMV promoter and expressed in COS or HEK293 cells. In this
30 activated constitutive state, antagonists of the receptor can be identified in the absence of

ligand by measuring the ability of a test compound to inhibit constitutive cell signaling activity. Appropriate assays for this are, again, cAMP assays, reporter gene activation assays or assays measuring the binding of GTP-gamma-S.

- 5 One preferred cell signaling assay is based upon the observation that cells stably transfected with DRRs show a change in intracellular calcium levels in response to incubation in the presence of angiotensin II or III (see Example 5). Thus, a procedure can be used to identify DRR agonists or antagonists that is similar to the radioreceptor assays discussed above except that angiotensin II or III is used instead of a labeled ligand and
- 10 calcium concentration is measured instead of bound radioactivity. The concentration of calcium in the presence of test compound and angiotensin II or III is compared with that in the presence of angiotensin II or III alone to determine whether the test compound is interacting at the DRR receptor. A statistically significant increase in intracellular calcium in response to test compound indicates that the test compound is acting as an agonist
- 15 whereas a statistically significant decrease in intracellular calcium indicates that it is acting as an antagonist.

V. Assay for Ability to Modulate DRR Expression

- One way to either increase or decrease the biological effects of a DRR is to alter the
- 20 extent to which the receptor is expressed in cells. Therefore, assays for the identification of compounds that either inhibit or enhance expression are of considerable interest. These assays are carried out by growing cells expressing a DRR in the presence of a test compound and then comparing receptor expression in these cells with expression in cells grown under essentially identical conditions but in the absence of the test compound. As in
- 25 the binding assays discussed above, it is desirable that the cells used be substantially free of competing G protein-coupled receptors. One way to quantitate receptor expression is to fuse the DRR sequence to a sequence encoding a peptide or protein that can be readily quantitated. For example, the DRR sequence may be ligated to a sequence encoding haemagglutinin as described in Example 5 and used to stably transfect cells. After

incubation with test compound the hemagglutinin/receptor complex can be immunoprecipitated and western blotted with anti- haemagglutinin antibody. Alternatively, Scatchard analysis of binding assays may be performed with labeled ligand to determine receptor number. The binding assays may be carried out as discussed above
5 and will preferably utilize cells that have been engineered to recombinantly express DRR.

A preferred group of test compounds for inclusion in the DRR expression assay consists of oligonucleotides complementary to various segments of the DRR nucleic acid sequence. These oligonucleotides should be at least 15 bases in length and should be derived from
10 non-conserved regions of the receptor nucleic acid sequence. Sequences may be based upon those shown as SEQ ID numbers 2, 4, 6, 8, 10, 12 or 14.

Oligonucleotides which are found to reduce receptor expression may be derivatized or conjugated in order to increase their effectiveness. For example, nucleoside
15 phosphorothioates may be substituted for their natural counterparts (see Cohen, J., Oligodeoxynucleotides, Antisense Inhibitors of Gene Expression, CRC Press (1989)). The oligonucleotides may be delivered to a patient in vivo for the purpose of inhibiting DRR expression. When this is done, it is preferred that the oligonucleotide be administered in a form that enhances its uptake by cells. For example, the oligonucleotide may be delivered
20 by means of a liposome or conjugated to a peptide that is ingested by cells (see e.g., U.S. Patent Nos. 4,897,355 and 4,394,448; see also non-U.S. patent documents WO 8903849 and EP 0263740). Other methods for enhancing the efficiency of oligonucleotide delivery are well known in the art and are also compatible with the present invention.

25 Having now described the invention, the same will be more readily understood through reference to the following Examples which are provided by way of illustration and which are not intended to limit the scope of the invention.

EXAMPLES

Example 1: Cloning of Rat DRR-1

Isolation of cDNA fragment.

5

Degenerate oligonucleotides were synthesized to highly conserved regions of G-protein coupled receptors (transmembrane spanning domains 2 and 7) with the following nucleotide sequences:

10 5' GG CCG TCG ACT TCA TCG TC(A/T) (A/C)(T/C)C TI(G/T) CI(T/C) TIG
C(A/C/G/T)G 3' (TM2:sense) SEQ ID NO:15; and

5' (A/G)(C/A/T)(A/T) (A/G)CA (A/G)TA IAT IAT IGG (A/G)TT 3'
(TM7:antisense) SEQ ID NO:16.

15

Poly A+ mRNA was isolated from cultured fetal rat dorsal root ganglia (Sprague-Dawley). The mRNA was reverse transcribed using the First Strand cDNA Synthesis kit (Pharmacia Biotech), subjected to an amplification reaction by polymerase chain reaction (PCR) using Ampli-Taq DNA (Perkin-Elmer Cetus) polymerase under the
20 following conditions: 3 minutes at 94 °C, 40 cycles of 1 minute at 94 °C, 45°C and 72 °C. A cDNA PCR fragment corresponding to approximately 650 bps was isolated and subcloned in pGEM-T-vector (Promega Corporation). The nucleotide sequence of the recombinant clone was determined using the T7-dideoxy chain termination sequencing kit (Pharmacia Biotech) and was found to be unique based upon searches of Genbank/EMBL
25 databases.

The full length rat DRR-1 sequence was obtained from rat genomic DNA using the 650 base pair fragment and the "Promoter Finder DNA Walking kit" (Clontech, cat # K1806-1). This kit contains five libraries of uncloned, adaptor-ligated genomic DNA fragments. The
30 procedure involves two consecutive PCR reactions. Both reactions were done using the

"Advantage Tth Polymerase Mix" also obtained from Clontech, following the conditions recommended by the vendor. The first PCR reaction was performed with the outer adaptor primer (AP1) provided in the kit and an outer, gene-specific primer (GSP1) derived from the sequence of the DRR-1 PCR fragment. The primary PCR mixtures were diluted and
5 used as a template for the secondary (nested) PCR reaction with the nested adapter primer (AP2) and a nested gene specific primer (GSP2). To obtain the sequence of the rat DRR-1 gene upstream of the sequence of the original PCR fragment, the following oligonucleotides were used:

10 GSP1: oligo YF3B59-B, 5'-CGCAGATGAGGTAGTACAGCATCAC SEQ ID NO:17
GSP2: oligo MML-R1, 5'-CTGTGAGAGAGATGGTAACATACAG SEQ ID NO:18

From the first library, a fragment AP2-MMLR1 of 1.9 Kb was obtained and from the third library, a fragment of approximately 1.0 Kb was obtained. To identify the sequence
15 downstream of the known sequence, the following primers were used:

GSP1: oligo YF3B59-F2, 5'-GCATCCTTGACTGGTTCTTCTCAG SEQ ID NO:19
GSP2: oligo MML-F1, 5'-GGGTGAGACTCATCATCATTTGTGG. SEQ ID NO:20

20 A fragment MMLF1-AP2 of approximately 1 Kb was obtained from the first library and a fragment of about 600 bp was obtained from the third library. The composite sequence of 1154 nucleotides containing the complete predicted open reading frame of DRR-1 is shown in Figure 1. The open reading frame codes for a 337 amino acid protein (Figure 2) with a predicted molecular mass of 38.7 kD. The protein sequence contains all the characteristic
25 features of G protein-coupled receptors: seven hydrophobic helices likely to represent transmembrane domains, potential glycosylation site at the N-terminal extracellular domain (position 30) and a conserved NPXXY sequence at position 285-289.

Example 2: Cloning of Human DRR Receptor Genes

Two approaches were used to identify and clone novel human DNA sequences homologous and/or related to the rat DRR-1 gene. First, a human genomic library was
5 screened in the lambda vector, Fix II, (Stratagene Cat.# 946203). Approximately 106 human genomic clones were plated and transferred onto nitrocellulose membranes for hybridization with the full length, 32P labeled, rat DRR-1 sequence as a probe. The hybridization was performed at 42 °C, overnight. The filters were washed at room temperature at low stringency (1X SSC/0.1% SDS) to allow detection of related but not
10 necessarily identical sequences.

The inserted human DNA present in positive phages was amplified by PCR using the "Expand PCR kit" from Boehringer-Mannheim under conditions allowing accurate amplification of very large fragments of DNA. These long fragments of DNA were
15 digested with various restriction enzymes and subcloned into a plasmid vector. The portions of these clones which hybridized with the rat DRR-1 gene probe were sequenced using the ABI cycle sequencing kit.

A second approach to identifying novel human sequences related to DRR-1 involved the
20 use of the polymerase chain reaction (PCR), performed on total human genomic DNA. Primers were synthesized based upon the human genomic clones described above and were as follows:

HML.H, 5'-GCAAGCTTTCTGAGCATGGATCCAACCGTC, SEQ ID 21 and
25 HML.Bg, 5'-CCCTCAGATCTCCAATTTGCTTCCCGACAG, SEQ ID NO:22.

Amplification resulted in a fragments of approximately 1 kilobase containing the entire coding sequence of the human genes. These fragments obtained were subcloned into the pGEM-T (Promega) vector for DNA sequencing analysis.

30

Using the above strategies, six human clones were isolated:

clone 7, SEQ ID numbers 3 and 4;

clone 18, SEQ ID numbers 5 and 6;

clone 23, SEQ ID numbers 7 and 8;

5 clone 24, SEQ ID numbers 9 and 10;

clone 36, SEQ ID numbers 11 and 12; and

clone 40, SEQ ID numbers 13 and 14.

None of these clones contain introns and their alignment may be seen in Figure 3.

10

At the amino acid sequence level, the rat DRR-1 clone is 47% to 49% identical to the human clones.

At the nucleic acid level, the rat DRR-1 clone is 56% to 58% identical to the human
15 clones. The level of sequence identity within the human clones (7, 18, 23, 24, 36, 40) is
very high, between 77% and 98% at the amino acid sequence level. All the human
sequences were used as queries to search for homologies in public databases (Genbank,
Swissprot, EST). No identical sequences were detected. The closest matches were to
members of the mas oncogene family of proteins. The overall amino acid sequence
20 homology between rat DRR-1 and any of the isolated human genes varied from 47 to 50 %.
However some stretches display a much higher level of sequence homology, particularly
the regions encoding the putative transmembrane domain III and VII (TM3 and TM7) and
the intracellular loops 2 and 3 where the homology between the rat sequence and its human
homologue is around 80%.

25

Example 3: In Situ Hybridization Experiments

Preparation of Tissue: Adult male Sprague-Dawley rats (~300 gm; Charles River,
St-Constant, Quebec) were sacrificed by decapitation. Brain and spinal cord with dorsal
root ganglia attached were removed, snap-frozen in isopentane at -40°C for 20 s and stored
30 at -80 °C. Frozen human brain, spinal cord and dorsal root ganglia were obtained from the

Brain and Tissue Bank for Developmental Disorders, University of Maryland at Baltimore, according to the strictest ethical guidelines. Frozen tissue was sectioned at 14 μ m in a Microm HM 500 M cryostat (Germany) and thaw-mounted onto ProbeOn Plus slides (Fisher Scientific, Montreal, Quebec). Sections were stored at -80°C prior to in situ hybridization.

Synthesis of Riboprobes: The plasmid pGemT-3b32 GPCR was linearized using either SacII and Not I restriction enzymes. Sense and antisense DRR riboprobes were transcribed in vitro using either T7 or SP6 RNA polymerases (Pharmacia Biotech), respectively in the presence of [35S]UTP (~800 Ci/mmol; Amersham, Oakville, Ontario). The plasmid pGemT-Clone 36 GPCR was linearized using SacII and Pst I restriction enzymes. Sense and antisense Clon36 riboprobes were transcribed in vitro using either SP6 or T7 RNA polymerases (Pharmacia Biotech), respectively in the presence of [35S]UTP. Following transcription, the DNA template was digested with DNase I (Pharmacia). Riboprobes were purified by phenol/chloroform/isoamyl alcohol extraction and precipitated in 70% ethanol containing ammonium acetate and tRNA. Quality of labeled riboprobes was verified by polyacrylamide-urea gel electrophoresis.

In situ Hybridization: Sections were postfixed in 4% paraformaldehyde (BDH, Poole, England) in 0.1 M phosphate buffer (pH 7.4) for 10 min at room temperature (RT) and rinsed in three changes of 2X standard sodium citrate buffer (SSC; 0.15 M NaCl, 0.015 M sodium citrate, pH 7.0). Sections were then equilibrated in 0.1 M triethanolamine, treated with 0.25% acetic anhydride in triethanolamine, rinsed in 2X SSC and dehydrated in an ethanol series (50-100%). Hybridization was performed in a buffer containing 75% formamide (Sigma, St-Louis, Mo), 600 mM NaCl, 10 mM Tris (pH 7.5), 1 mM EDTA, 1X Denhardt's solution (Sigma), 50 (g/ml) denatured salmon sperm DNA (Sigma), 50 (g/ml) yeast tRNA (Sigma), 10% dextran sulfate (Sigma), 20 mM dithiothreitol and [35S]UTP-labeled cRNA probes (10 X10⁶ cpm/ml) at 55°C for 18 h in humidified chambers. Following hybridization, slides were rinsed in 2X SSC at RT, treated with 20 (g/ml) RNase IA (Pharmacia) in RNase buffer (10 mM Tris, 500 mM NaCl, 1 mM EDTA, pH 7.5) for 45

min at RT and washed to a final stringency of 0.1X SSC at 65 °C. Sections were then dehydrated and exposed to Kodak Biomax MR film for 21 days and/or dipped in Kodak NTB2 emulsion diluted 1:1 with distilled water and exposed for 4-6 weeks at 4°C prior to development and counterstaining with cresyl violet acetate (Sigma).

5

Results: Of all regions examined within the neuraxis of the rat, DRR-1 mRNA was exclusively expressed in dorsal root ganglia. High resolution emulsion autoradiography showed accumulations of silver grains exclusively over small and some medium size neurons. This unique and highly restricted distribution pattern for DRR-1 was confirmed in the rat embryo. Sagittal section of an E17 rat fetus showed that DRR-1 mRNA is confined to DRGs. All other structures of the rat embryo were devoid of any specific hybridization signal reinforcing the highly selective nature of DRR-1 expression

The expression of human Clone 36 receptor was present in human fetal dorsal root ganglia but not in spinal cord. Specific hybridization signal for Clone 36 was not detected in any of the human adult CNS tissues examined thus far. These include spinal cord, cortex, hippocampus, thalamus, substantia nigra and periaqueductal gray (data not shown). Presence of Clone 36 mRNA in adult DRGs remains to be examined. Standard controls in which additional spinal cord with DRG sections were hybridized with rat DRR-1 antisense or Clone 36 sense 35S-labeled probes displayed no specific hybridization signal.

Example 4: Northern Blots

Commercial rat and human multiple Northern blots containing 2 g of polyA RNA from various tissues (Clontech) were used to determine the expression and distribution of the rat DRR-1 message and its human homologues. Radioactively labeled probes were prepared as follows: twenty five ng of a 650 bp 3b-32 PCR fragment derived from rat DRR-1 (see Example 1) or human clone 36 were random-prime labeled using the Ready-to-Go DNA labeling kit (Pharmacia Biotech) and [32P]CTP (3000 Ci/mmol/Amersham). The blot was prehybridized for 1 hour at 68 °C using Expresshyb (Clontech) followed by hybridization (2X10⁶ cpm/ml of probe) for one hour using the same conditions. Blots were washed at

30

room temperature in 2X SSC, 0.05% SDS for 30 min. followed by 3x washes in 0.2X SSC, 1 % SDS at 50 °C for 60 min. and exposed at -80 °C to Kodak Biomax film for 6 days.

- 5 **Expression and Distribution of rat DRR-1:** All the rat tissues studied (heart, brain, spleen, lung, skeletal muscle, kidney and testis) were negative for the expression of DRR-1 following 2 weeks exposure whereas rat genomic Southern analysis revealed a 1.1 kb band when probed with the same cDNA fragment.
- 10 **Expression and Distribution of Human Clone 36:** Northern blots containing RNA from various human tissues were probed with a radio-labeled DNA fragment from clone 36. All the human tissues studied (human fetal brain, lung, liver and kidney and adult human cerebellum, cerebral cortex, medulla, occipital pole, frontal lobe, temporal lobe, putamen, spinal cord, amygdala, caudate nucleus, corpus callosum, hippocampus, total brain,
- 15 subthalamic nucleus and thalamus) were negative for the expression of this receptor following 2 weeks exposure.

Example 5: Calcium Signaling in Response to Angiotensin I-III

The coding sequence of human clone 24 was transferred into a pcDNA3 vector and
20 modified to add a haemagglutinin tag at the C-terminus of the receptor sequence. This clone, designated as pcDNA3-HML-HA24 was transfected into HEK293 cells using a modified CaCl₂ method (Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press (1989)). The cells were maintained in culture medium at 37 °C, 5% CO₂ and diluted 10 fold every 3 days.

25 The cells were inoculated in 90 mm tissue culture dishes (5 x 10⁵ cells per flask) in Dulbecco's Modified Essential Medium (DMEM, Gibco BRL), supplemented with 10% fetal bovine serum (FBS), 100 U/ml penicillin, 100 µg/ml streptomycin and 0.25 µg/ml fungizone. One day after inoculation, cells were transiently transfected with 30 µg of
30 plasmid DNA per dish. The cells were harvested 48 hours post transfection for analysis.

The expression of the gene was first checked by immunoprecipitation and western blotting with an anti-haemagglutinin antibody. A protein of approximately 43 KD was detected in stably as well as transiently transfected HEK293 cells, but not in control cells.

5 Stably transfected HEK293 cells were obtained after approximately 21 days of selection in culture medium containing 800 µg/ml G418. Calcium signaling measurement was performed with one of these stably transfected cell line, 293/pcDNA3-HML-HA24. The cells were grown on a 24 mm round glass cover slides to 50-70% confluence. After rinsing the cells with 1.8 NBS buffer (135 mM NaCl, 5 mM KCl, 1.2 mM MgCl₂, 1.8 mM
10 CaCl₂, 5 mM glucose and 10 mM HEPES, pH 7.3), the cells were incubated for one hour at room temperature in the presence of 0.5 ml of 3.5 µM FURA-2 AM (Molecular Probe, F-1221) diluted in 1.8 NBS. The cells were then rinsed three times with 1.8 NBS and incubated for a further 30 minutes at room temperature. The calcium displacement was measured using a PTI (Photon Technology International) D104 photometer equipped with a
15 PTI Delta RAM High speed multiwavelength illuminator, a PTI SC500 Shutter controller, a PTI LPS220 ARC lamp supply and the PTI FELIX software, v.1.2. Groups of 2 to 8 cells were chosen and isolated with the photometer diaphragm. The cells were exposed to 340 and 380 nm light and the 510 nm light emitted by the cells was recorded. Angiotensin I, II and III, were added successively - in various order from one experiment to the next -
20 followed by bradykinin as a positive control. Upon stimulation with angiotensin II and angiotensin III, a significant response was obtained. Addition of angiotensin I produced no response.

All references cited herein are fully incorporated by reference. Having now fully
25 described the invention, it will be understood by one of skill in the art that the invention may be performed within a wide and equivalent range of conditions, parameters and the like, without affecting the spirit or scope of the invention or any embodiment thereof.

Applicant's or agent's file reference	N 1807-1 WO	International application No.
--	-------------	-------------------------------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>6</u> line <u>8-9</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ)	
Address of depositary institution (including postal code and country) Mascheroder Weg 1b D-38124 Braunschweig Germany	
Date of deposit 27 November 1997	Accession Number DSM 11877
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
In respect of all designated states in which such action is possible and to the extent that it is legally permissible under the law of the designated state, it is requested that a sample of the deposited micro-organism be made available only by the issue thereof to an independent expert, in accordance with the relevant patent legislation, e.g. Rule 28(4) EPC, and generally similar provisions <i>mutatis mutandis</i> for any other designated state.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

SUBSTITUTE SHEET (RULE 26)

CLAIMS

1. A protein, except as existing in nature, comprising the amino acid sequence consisting functionally of rat dorsal root receptor 1 (DRR-1) as shown in SEQ ID NO:1.
5
2. A substantially pure protein according to claim 1, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:1.
3. A substantially pure polynucleotide encoding a protein comprising the amino acid
10 sequence consisting functionally of rat DRR-1 as shown in SEQ ID NO:1.
4. The polynucleotide of claim 3, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:1.
- 15 5. The polynucleotide of claim 4, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:2.
6. A protein, except as existing in nature, comprising the amino acid sequence consisting functionally of human dorsal root receptor 1 (DRR-1) as shown in SEQ ID NO:3.
20
7. A substantially pure protein according to claim 6, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:3.
8. A substantially pure polynucleotide encoding a protein comprising the amino acid
25 sequence consisting functionally of human DRR-1 as shown in SEQ ID NO:3.
9. The polynucleotide of claim 8, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:3.

10. The polynucleotide of claim 9, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:4.
11. A protein, except as existing in nature, comprising the amino acid sequence consisting
5 functionally of human dorsal root receptor 2 (DRR-2) as shown in SEQ ID NO:5.
12. A substantially pure protein according to claim 11, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:5.
- 10 13. A substantially pure polynucleotide encoding a protein comprising the amino acid sequence consisting functionally of human DRR-2 as shown in SEQ ID NO:5.
14. The polynucleotide of claim 13, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:5.
- 15 15. The polynucleotide of claim 14, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:6.
16. A protein, except as existing in nature, comprising the amino acid sequence consisting
20 functionally of human dorsal root receptor 3 (DRR-3) as shown in SEQ ID NO:7.
17. A substantially pure protein according to claim 16, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:7.
- 25 18. A substantially pure polynucleotide encoding a protein comprising the amino acid sequence consisting functionally of human DRR-3 as shown in SEQ ID NO:7.
19. The polynucleotide of claim 18, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:7.

20. The polynucleotide of claim 19, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:8.
21. A protein, except as existing in nature, comprising the amino acid sequence consisting functionally of human dorsal root receptor 4 (DRR-4) as shown in SEQ ID NO:9.
22. A substantially pure protein according to claim 21, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:9.
23. A substantially pure polynucleotide encoding a protein comprising the amino acid sequence consisting functionally of human DRR-4 as shown in SEQ ID NO:9.
24. The polynucleotide of claim 23, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:9.
25. The polynucleotide of claim 24, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:10.
26. A protein, except as existing in nature, comprising the amino acid sequence consisting functionally of human dorsal root receptor 5 (DRR-5) as shown in SEQ ID NO:11.
27. A substantially pure protein according to claim 26, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:11.
28. A substantially pure polynucleotide encoding a protein comprising the amino acid sequence consisting functionally of human DRR-5 as shown in SEQ ID NO:11.
29. The polynucleotide of claim 28, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:11.

30. The polynucleotide of claim 29, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:12.
31. A protein, except as existing in nature, comprising the amino acid sequence consisting
5 functionally of human dorsal root receptor 6 (DRR-6) as shown in SEQ ID NO:13.
32. A substantially pure protein according to claim 31, wherein said amino acid sequence consists essentially of the amino acid sequence of SEQ ID NO:13.
- 10 33. A substantially pure polynucleotide encoding a protein comprising the amino acid sequence consisting functionally of human DRR-6 as shown in SEQ ID NO:13.
34. The polynucleotide of claim 33, wherein said polynucleotide encodes a protein consisting essentially of the amino acid sequence of SEQ ID NO:13.
- 15 35. The polynucleotide of claim 34, wherein said polynucleotide has the nucleotide sequence of SEQ ID NO:14.
36. An antibody made by a process comprising the step of injecting a pharmaceutically
20 acceptable preparation comprising the protein of anyone of claims 1, 2, 6, 7, 11, 12, 16, 17, 21, 22, 26, 27, 31 or 32, into an animal capable of producing said antibody.
37. An antibody that binds specifically to anyone of the proteins of claims 1, 2, 6, 7, 11, 12, 16, 17, 21, 22, 26, 27, 31 or 32
- 25 38. A vector for expressing rat DRR-1, comprising the polynucleotide of either one of claim 3 or 4.

39. A vector for expressing anyone of
- (i) human DRR-1, comprising a polynucleotide of claim 8 or 9;
 - (ii) human DRR-2, comprising a polynucleotide of claim 13 or 14;
 - 5 (iii) human DRR-3, comprising a polynucleotide of claim 18 or 19;
 - (iv) human DRR-4, comprising a polynucleotide of claim 23 or 24;
 - (v) human DRR-5, comprising a polynucleotide of claim 28 or 29;
 - (vi) human DRR-6, comprising a polynucleotide of claim 33 or 34.
- 10 40. A host cell transformed with a vector according to claim 38 or 39.
41. Recombinant rat DRR-1, human DRR-1, human DRR-2, human DRR-3,
human DRR-4, human DRR-5, human DRR-6, produced by the host cell of claim 40.
- 15 42. A method for assaying a test compound for its ability to bind or to activate a G
protein-coupled dorsal root ganglia specific receptor (DRR), comprising:
- a) incubating a source containing DRR but substantially free of other G protein-
coupled receptors, with
 - i) a ligand known to bind to DRR;
 - 20 ii) said test compound; and
 - b) determining the extent to which said ligand binding is displaced by said test
compound.

25

30

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT: Astra Pharma Inc. Canada

(ii) TITLE OF INVENTION: Novel receptor

10 (iii) NUMBER OF SEQUENCES: 22

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Astra AB, Patent Department

(B) STREET: S-151 85 Södertälje

15 (C) CITY: Södertälje

(D) STATE:

(E) COUNTRY: Sweden

(F) ZIP: none

20 (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

30

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 46-8 553 26000

(B) TELEFAX: 46-8 553 28820

5

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 337 amino acids

10 (B) TYPE: amino acid

(C) STRANDEDNESS: not relevant

(D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

15

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Met Val Cys Val Leu Arg Asp Thr Thr Gly Arg Phe Val Ser Met Asp

1 5 10 15

25 Pro Thr Ile Ser Ser Leu Ser Thr Glu Ser Thr Thr Leu Asn Lys Thr

20 25 30

Gly His Pro Ser Cys Arg Pro Ile Leu Thr Leu Ser Phe Leu Val Pro

35 40 45

30

Ile Ile Thr Leu Leu Gly Leu Ala Gly Asn Thr Ile Val Leu Trp Leu
50 55 60

Leu Gly Phe Arg Met Arg Arg Lys Ala Ile Ser Val Tyr Val Leu Asn
5 65 70 75 80

Leu Ser Leu Ala Asp Ser Phe Phe Leu Cys Cys His Phe Ile Asp Ser
85 90 95

Leu Met Arg Ile Met Asn Phe Tyr Gly Ile Tyr Ala His Lys Leu Ser
10 100 105 110

Lys Glu Ile Leu Gly Asn Val Ala Phe Ile Pro Tyr Ile Ser Gly Leu
115 120 125

Ser Ile Leu Ser Ala Ile Ser Thr Glu Arg Cys Leu Ser Val Leu Trp
130 135 140

Pro Ile Trp Tyr His Cys His Arg Pro Arg Asn Met Ser Ala Ile Ile
145 150 155 160

Cys Val Leu Ile Trp Val Leu Ser Phe Leu Met Gly Ile Leu Asp Trp
165 170 175

Phe Phe Ser Gly Phe Leu Gly Glu Thr His His His Leu Trp Lys Asn
180 185 190

Val Asp Phe Ile Val Thr Ala Phe Leu Ile Phe Leu Phe Met Leu Leu
195 200 205

30

SUBSTITUTE SHEET (RULE 26)

Phe Gly Ser Ser Leu Ala Leu Leu Val Arg Ile Leu Cys Gly Ser Arg
210 215 220

Arg Lys Pro Leu Ser Arg Leu Tyr Val Thr Ile Ser Leu Thr Val Met
5 225 230 235 240

Val Tyr Leu Ile Cys Gly Leu Pro Leu Gly Leu Tyr Leu Phe Leu Leu
245 250 255

Tyr Trp Phe Gly Ile His Leu His Tyr Pro Phe Cys His Ile Tyr Gln
10 260 265 270

Val Thr Val Leu Leu Ser Cys Val Asn Ser Ser Ala Asn Pro Ile Ile
275 280 285

Tyr Phe Leu Val Gly Ser Phe Arg His Arg Lys Lys His Arg Ser Leu
15 290 295 300

Lys Met Val Leu Lys Arg Ala Leu Glu Glu Thr Pro Glu Glu Asp Glu
20 305 310 315 320

Tyr Thr Asp Ser His Val Gln Lys Pro Thr Glu Ile Ser Glu Arg Arg
325 330 335

Cys
25

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 1011 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ATGGTTTGTG TTCTCAGGGA CACTACTGGA AGATTGTGA GCATGGATCC AACCATCTCA 60
20 TCCCTCAGTA CAGAATCTAC AACACTGAAT AAAACTGGTC ATCCCAGTTG CAGGCCAATC 120
CTCACCCCTGT CCTTCCTGGT CCCCATCATC ACCCTGCTTG GATTGGCAGG AAACACCATT 180
GTACTCTGGC TCTTGGGATT CCGCATGCGC AGGAAAGCCA TCTCAGTCTA CGTCCTCAAC 240
25 CTGTCTCTGG CAGACTCCTT CTTCTCTGTC TGCCATTTTA TTGACTCTCT GATGCGGATC 300
ATGAACTTCT ATGGCATCTA TGCCCATAAA TTAAGCAAAG AAATCTTAGG CAATGTAGCA 360
30 TTCATTCCCT ATATCTCAGG CCTGAGCATC CTCAGTGCTA TCAGCACGGA GCGCTGCCTG 420

SUBSTITUTE SHEET (RULE 26)

TCTGTATTGT GGCCAATCTG GTACCACTGC CACCGCCCAA GAAACATGTC AGCTATTATA 480

TGTGTTCTAA TCTGGGTTCT GTCCTTTCTC ATGGGCATCC TTGACTGGTT TTTCTCAGGA 540

5 TCCCTGGGTG AGACTCACCA TCATTTGTGG AAAAATGTTG ACTTTATTGT AACTGCATTT 600

CTGATTTTTT TATTTATGCT TCTCTTTGGG TCCAGTCTGG CGCTACTGGT GAGGATCCTC 660

10 TGTGGTTCCA GACGGAAACC ACTGTCCAGG CTGTACGTTA CAATCTCTCT CACAGTGATG 720

GTCTACCTCA TCTGCGGCCT GCCTCTCGGG CTTTACTTGT TCCTGCTATA TTGGTTTGGG 780

ATCCATTAC ATTATCCCTT TTGTCACATT TACCAAGTTA CTGTGCTCCT GTCCTGTGTG 840

15 AACAGCTCTG CCAACCCCAT CATTTACTTC CTTGTAGGGT CCTTTAGGCA CCGTAAAAAG 900

CATCGGTCCC TCAAAATGGT TCTTAAAAGG GCTCTGGAGG AGACTCCTGA GGAGGATGAA 960

20 TATACAGACA GCCATGTTCA GAAACCCACT GAGATCTCAG AAAGGAGATG T 1011

(2) INFORMATION FOR SEQ ID NO:3:

- 25 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 322 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: not relevant

30

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

5 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met Asp Pro Thr Ile Pro Val Leu Gly Thr Lys Leu Thr Pro Ile Asn
10 1 5 10 15
Gly Arg Glu Glu Thr Pro Cys Tyr Asn Gln Thr Leu Ser Phe Thr Gly
20 25 30
Leu Thr Cys Ile Ile Ser Leu Val Ala Leu Thr Gly Asn Ala Val Val
15 35 40 45
Leu Trp Leu Leu Gly Cys Arg Met Arg Arg Asn Ala Val Ser Ile Tyr
20 50 55 60
Ile Leu Asn Leu Val Ala Ala Asn Phe Leu Phe Leu Ser Gly His Ile
65 70 75 80
Ile Phe Ser Pro Leu Pro Leu Ile Asn Ile Arg His Pro Ile Ser Lys
25 85 90 95
Ile Leu Ser Pro Val Met Thr Phe Pro Tyr Phe Ile Gly Leu Ser Met
100 105 110

Leu Ser Ala Ile Ser Thr Glu Arg Cys Leu Ser Ile Leu Trp Pro Ile
 115 120 125

5 Trp Tyr His Cys Arg Arg Pro Arg Tyr Leu Ser Ser Val Met Cys Val
 130 135 140

Leu Leu Trp Ala Leu Ser Leu Leu Arg Ser Ile Leu Glu Trp Met Phe
 145 150 155 160

10 Cys Asp Phe Leu Phe Ser Gly Ala Asn Ser Val Trp Cys Glu Thr Ser
 165 170 175

Asp Phe Ile Thr Ile Ala Trp Leu Val Phe Leu Cys Val Val Leu Cys
 15 180 185 190

Gly Ser Ser Leu Val Leu Leu Val Arg Ile Leu Cys Gly Ser Arg Lys
 195 200 205

20 Met Pro Leu Thr Arg Leu Tyr Val Thr Ile Leu Leu Thr Val Leu Val
 210 215 220

Phe Leu Leu Cys Gly Leu Pro Phe Gly Ile Gln Trp Ala Leu Phe Ser
 225 230 235 240

25 Arg Ile His Leu Asp Trp Lys Val Leu Phe Cys His Val His Leu Val
 245 250 255

Ser Ile Phe Leu Ser Ala Leu Asn Ser Ser Ala Asn Pro Ile Ile Tyr
 30 260 265 270

SUBSTITUTE SHEET (RULE 26)

Phe Phe Val Gly Ser Phe Arg Gln Arg Gln Asn Arg Gln Asn Leu Lys

275

280

285

5 Leu Val Leu Gln Arg Ala Leu Gln Asp Thr Pro Glu Val Asp Glu Gly

290

295

300

Gly Gly Trp Leu Pro Gln Glu Thr Leu Glu Leu Ser Gly Ser Lys Leu

10

305

310

315

320

Glu Gln

15 (2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 969 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

25 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

30

SUBSTITUTE SHEET (RULE 26)

ATGGATCCAA CCATCCCAGT CTTGGGTACA AAACTGACAC CAATCAACGG ACGTGAGGAG 60

ACTCCTTGCT ACAACCAAAC CCTGAGCTTC ACGGGGCTGA CGTGCATCAT TTCCCTTGTC 120

5 GCGCTGACAG GAAACGCGGT TGTGCTCTGG CTCCTGGGCT GCCGCATGCG CAGGAACGCT 180

GTCTCCATCT ACATCCTCAA CCTGGTCGCG GCCAACTTCC TCTTCCTTAG CGGCCACATT 240

ATATTTTCGC CGTTACCCCT CATCAATATC CGCCATCCCA TCTCCAAAAT CCTCAGTCCT 300

10 GTGATGACCT TTCCCTACTT TATAGGCCTA AGCATGCTGA GCGCCATCAG CACCGAGCGC 360

TGCCTGTCCA TCCTGTGGCC CATCTGGTAC CACTGCCGCC GCCCCAGATA CCTGTCATCG 420

15 GTCATGTGTG TCCTGCTCTG GGCCCTGTCC CTGCTGCGGA GTATCCTGGA GTGGATGTTT 480

TGTGACTTCC TGTTTAGTGG TGCTAATTCT GTTTGGTGTG AAACGTCAGA TTTCATTACA 540

ATCGCGTGGC TGGTTTTTTT ATGTGTGGTT CTCTGTGGGT CCAGCCTGGT CCTGCTGGTC 600

20 AGGATTCTCT GTGGATCCCG GAAGATGCCG CTGACCAGGC TGTACGTGAC CATCCTCCTC 660

ACAGTGCTGG TCTTCCTCCT CTGTGGCCTG CCCTTTGGCA TTCAGTGGGC CCTGTTTTCC 720

25 AGGATCCACC TGGATTGGAA AGTCTTATTT TGTATGTGTC ATCTAGTTTC CATTTTCCTG 780

TCCGCTCTTA ACAGCAGTGC CAACCCCATC ATTTACTTCT TCGTGGGCTC CTTTAGGCAG 840

CGTCAAAATA GGCAAAACCT GAAGCTGGTT CTCCAAAGGG CTCTGCAGGA CACGCCTGAG 900

30

GTGGATGAAG GTGGAGGGTG GCTTCCTCAG GAAACCTGG AGCTGTCGGG AAGCAAATTG 960

GAGCAGTGA 969

5

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 322 amino acids

10 (B) TYPE: amino acid

(C) STRANDEDNESS: not relevant

(D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

15

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Asp Pro Thr Val Pro Val Leu Gly Thr Glu Leu Thr Pro Ile Asn

1 5 10 15

25 Gly Arg Glu Glu Thr Pro Cys Tyr Lys Gln Thr Leu Ser Phe Thr Gly

20 25 30

Leu Thr Cys Ile Val Ser Leu Val Ala Leu Thr Gly Asn Ala Val Val

35 40 45

30

SUBSTITUTE SHEET (RULE 26)

Leu Trp Leu Leu Gly Cys Arg Met Arg Arg Asn Ala Val Ser Ile Tyr
 50 55 60

Ile Leu Asn Leu Val Ala Ala Asp Phe Leu Phe Leu Ser Gly His Ile
 5 65 70 75 80

Ile Cys Ser Pro Leu Arg Leu Ile Asn Ile Ser His Pro Ile Ser Lys
 85 90 95

10 Ile Leu Ser Pro Val Met Thr Phe Pro Tyr Phe Ile Gly Leu Ser Met
 100 105 110

Leu Asn Ala Ile Ser Thr Glu Arg Cys Leu Ser Ile Leu Trp Pro Ile
 115 120 125

15 Trp Tyr His Cys Arg Arg Pro Arg Tyr Leu Ser Ser Val Met Cys Val
 130 135 140

Leu Leu Trp Ala Pro Ser Leu Leu Arg Ser Ile Leu Glu Trp Met Phe
 20 145 150 155 160

Cys Asp Phe Leu Phe Ser Gly Ala Asp Ser Val Arg Cys Glu Thr Ser
 165 170 175

25 Asp Phe Ile Thr Ile Ala Trp Leu Val Phe Leu Arg Val Val Leu Cys
 180 185 190

Gly Ser Ser Leu Val Leu Leu Val Arg Ile Leu Cys Gly Ser Arg Lys
 195 200 205

30

Met Pro Leu Thr Arg Leu Tyr Val Thr Ile Leu Leu Thr Val Leu Val
 210 215 220

Phe Leu Leu Cys Gly Leu Pro Phe Gly Ile Gln Trp Ala Leu Phe Ser
 5 225 230 235 240

Arg Ile His Leu Asp Trp Lys Val Leu Phe Cys His Val His Leu Val
 245 250 255

10

Ser Ile Phe Leu Ser Ala Leu Asn Ser Ser Ala Asn Pro Ile Ile Tyr
 260 265 270

Phe Phe Met Gly Ser Phe Arg Gln Leu Gln Asn Arg Lys Thr Leu Lys
 15 275 280 285

Leu Val Leu Gln Arg Asp Leu Gln Asp Thr Pro Glu Val Asp Glu Gly
 290 295 300

Gly Trp Trp Leu Pro Gln Glu Thr Leu Glu Leu Ser Gly Ser Lys Leu
 20 305 310 315 320

Glu Ile

25

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 969 base pairs

30

(B) TYPE: nucleic acid

SUBSTITUTE SHEET (RULE 26)

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

ATGGATCCAA CCGTCCCAGT CTTGGGTACA GAACTGACAC CAATCAACGG ACGTGAGGAG 60

ACTCCTTGCT ACAAGCAGAC CCTGAGCTTC ACGGGGCTGA CGTGCATCGT TTCCCTTGTC 120

15

GCGCTGACAG GAAACGCGGT TGTGCTCTGG CTCCTGGGCT GCCGCATGCG CAGGAACGCT 180

GTCTCCATCT ACATCCTCAA CCTGGTCCGG GCCGACTTCC TCTTCCTTAG CGGCCACATT 240

20

ATATGTTTCGC CGTTACGCCT CATCAATATC AGCCATCCCA TCTCCAAAAT CCTCAGTCCT 300

GTGATGACCT TTCCCTACTT TATAGGCCTA AGCATGCTGA ACGCCATCAG CACCGAGCGC 360

TGCCTGTCCA TCCTGTGGCC CATCTGGTAC CACTGCCGCC GCCCCAGATA CCTGTCATCG 420

25

GTGATGTGTG TCCTGCTCTG GGCCCCGTCC CTGCTGCGGA GTATCCTGGA GTGGATGTTT 480

TGTGACTTCC TGTTTAGTGG TGCTGATTCT GTTCGGTGTG AAACGTCAGA TTTCATTACA 540

30

ATCGCGTGGC TGGTTTTTTT ACGTGTGGTT CTCTGTGGGT CCAGCCTGGT CCTGCTGGTC 600

SUBSTITUTE SHEET (RULE 26)

AGGATTCTCT GTGGATCCCG GAAGATGCCG CTGACCAGGC TGTACGTGAC CATCCTCCTC 660

ACAGTGCTGG TCTTCCTCCT CTGTGGCCTG CCCTTTGGCA TTCAGTGGGC CCTGTTTTC 720

5 AGGATCCACC TGGATTGGAA AGTCTTATTT TGTCAIGTGC ATCTAGTTTC CATTTCCTG 780

TCCGCTCTTA ACAGCAGTGC CAACCCCATC ATTTACTTCT TCATGGGCTC CTTTAGGCAG 840

10 CTTCAAAACA GGAAGACCCT CAAGCTGGTT CTCCAGAGGG ATCTGCAGGA CACGCCTGAG 900

GTGGATGAAG GTGGATGGTG GCTTCCTCAG GAAACCCTGG AGCTGTCGGG AAGCAAATTG 960

GAGATCTGA 969

15

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 322 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: not relevant

25 (ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

30

SUBSTITUTE SHEET (RULE 26)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

	Met	Asp	Pro	Thr	Val	Ser	Thr	Leu	Asp	Thr	Glu	Leu	Thr	Pro	Ile	Asn	
	1					5					10				15		
5																	
	Gly	Thr	Glu	Glu	Thr	Leu	Cys	Tyr	Lys	Gln	Thr	Leu	Ser	Leu	Thr	Val	
						20				25					30		
	Leu	Thr	Cys	Ile	Val	Ser	Leu	Val	Gly	Leu	Thr	Gly	Asn	Ala	Val	Val	
10						35				40					45		
	Leu	Trp	Leu	Leu	Gly	Cys	Arg	Met	Arg	Arg	Asn	Ala	Phe	Ser	Ile	Tyr	
						50				55					60		
15	Ile	Leu	Asn	Leu	Ala	Ala	Ala	Asp	Phe	Leu	Phe	Leu	Ser	Gly	Arg	Leu	
	65						70					75				80	
	Ile	Tyr	Ser	Leu	Leu	Ser	Phe	Ile	Ser	Ile	Pro	His	Thr	Ile	Ser	Lys	
							85				90				95		
20																	
	Ile	Leu	Tyr	Pro	Val	Met	Met	Phe	Ser	Tyr	Phe	Ala	Gly	Leu	Asn	Phe	
							100				105				110		
	Leu	Ser	Ala	Val	Ser	Thr	Asp	Arg	Cys	Leu	Ser	Val	Leu	Trp	Pro	Ile	
25							115				120				125		
	Trp	Tyr	Arg	Cys	His	Arg	Pro	Thr	His	Leu	Ser	Ala	Val	Val	Cys	Val	
							130				135				140		

SUBSTITUTE SHEET (RULE 26)

Leu Leu Trp Ala Leu Ser Leu Leu Arg Ser Ile Leu Glu Trp Met Leu
 145 150 155 160

5 Cys Gly Phe Leu Phe Ser Gly Ala Asp Ser Ala Trp Cys Gln Thr Ser
 165 170 175

Asp Phe Ile Thr Val Ala Trp Leu Ile Phe Leu Cys Val Val Leu Cys
 180 185 190

10 Gly Ser Ser Leu Val Leu Leu Ile Arg Ile Leu Cys Gly Ser Arg Lys
 195 200 205

15 Ile Pro Leu Thr Arg Leu Tyr Val Thr Ile Leu Leu Thr Val Leu Val
 210 215 220

Phe Leu Leu Cys Gly Leu Pro Phe Gly Ile Gln Phe Phe Leu Phe Leu
 225 230 235 240

20 Trp Ile His Val Asp Arg Glu Val Leu Phe Cys His Val His Leu Val
 245 250 255

Ser Ile Phe Leu Ser Ala Leu Asn Ser Ser Ala Asn Pro Ile Ile Tyr
 25 260 265 270

Phe Phe Val Gly Ser Leu Arg Gln Arg Gln Asn Arg Gln Asn Leu Lys
 275 280 285

SUBSTITUTE SHEET (RULE 26)

Leu Val Leu Gln Arg Ala Leu Gln Asp Thr Pro Glu Val Asp Glu Gly

290

295

300

5

Gly Gly Trp Leu Pro Gln Glu Thr Leu Glu Leu Ser Gly Ser Arg Leu

305

310

315

320

Glu In

10

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 969 base pairs

15

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

20

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

ATGGATCCAA CCGTCTCAAC CTTGGACACA GAACTGACAC CAATCAACGG AACTGAGGAG 60

ACTCTTTGCT ACAAGCAGAC CTTGAGCCTC ACGGTGCTGA CGTGATCGT TTCCCTTGTC 120

30

SUBSTITUTE SHEET (RULE 26)

GGGCTGACAG GAAACCCAGT TGTACTGTGG CTCTGGGGCT GCCGCATGCG CAGGAACGCC 180

TTCTCCATCT ACATCCTCAA CTGGCCGCA GCAGACTTCC TCTTCCTCAG CGGCCGCCTT 240

5 ATATATTCCC TGTTAAGCTT CATCAGTATC CCCCATACCA TCTCTAAAAT CCTCTATCCT 300

GTGATGATGT TTTCTACTT TGCAGGCCTG AACTTTCTGA GTGCCGTGAG CACCGATCGC 360

TGCCTGTCCG TCCTGTGGCC CATCTGGTAC CGCTGCCACC GCCCCACACA CCTGTCAGCG 420

10 GTGGTGTGTG TCCTGCTCTG GGCCCTGTCC CTGCTGCGGA GCATCCTGGA ATGGATGTTA 480

TGTGGCTTCC TGTTCAGTGG TGCTGATTCT GCTTGGTGTC AAACATCAGA TTTATCACA 540

15 GTCGCGTGGC TGATTTTTTT ATGTGTGGTT CTCTGTGGGT CCAGCCTGGT CCTGCTGATC 600

AGGATTCTCT GTGGATCCCG GAAGATACCG CTGACCAGGC TGTACGTGAC CATCCTGCTC 660

ACAGTACTGG TCTTCCTCCT CTGTGGCCTG CCCTTTGGCA TTCAGTTTTT CCTATTTTTA 720

20 TGGATCCACG TGGACAGGGA AGTCTTATTT TGTCATGTGC ATCTAGTTTC CATTTTCCTG 780

TCCGCTCTTA ACAGCAGTGC CAACCCCATC ATTTACTTCT TCGTGGGCTC CCTTAGGCAG 840

25 CGTCAAAATA GGCAGAACCT GAAGCTGGTT CTCCAGAGGG CTCTGCAGGA CACGCCTGAG 900

GTGGATGAAG GTGGAGGGTG GCTTCCTCAG GAAACCCTGG AGCTGTCGGG AAGCAGATTG 960

GAGCAGTGA 969

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 322 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: not relevant

10 (ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Met Asp Pro Thr Val Ser Thr Leu Asp Thr Glu Leu Thr Pro Ile Asn

1 5 10 15

20

Gly Thr Glu Glu Thr Leu Cys Tyr Lys Gln Thr Leu Ser Leu Thr Val

20 25 30

Leu Thr Cys Ile Val Ser Leu Val Gly Leu Thr Gly Asn Ala Val Val

25 35 40 45

Leu Trp Leu Leu Gly Cys Arg Met Arg Arg Asn Ala Phe Ser Ile Tyr

50 55 60

SUBSTITUTE SHEET (RULE 26)

Ile Leu Asn Leu Ala Ala Ala Asp Phe Leu Phe Leu Ser Gly Arg Leu
 65 70 75 80

5 Ile Tyr Ser Leu Leu Ser Phe Ile Ser Ile Pro His Thr Ile Ser Lys
 85 90 95

Ile Leu Tyr Pro Val Met Met Phe Ser Tyr Phe Ala Gly Leu Ser Phe
 100 105 110

10 Leu Ser Ala Val Ser Thr Glu Arg Cys Leu Ser Val Leu Trp Pro Ile
 115 120 125

Trp Tyr Arg Cys His Arg Pro Thr His Leu Ser Ala Val Val Cys Val
 15 130 135 140

Leu Leu Trp Ala Leu Ser Leu Leu Arg Ser Ile Leu Glu Trp Met Leu
 145 150 155 160

20 Cys Gly Phe Leu Phe Ser Gly Ala Asp Ser Ala Trp Cys Gln Thr Ser
 165 170 175

Asp Phe Ile Thr Val Ala Trp Leu Ile Phe Leu Cys Val Val Leu Cys
 180 185 190

25 Gly Ser Ser Leu Val Leu Leu Ile Arg Ile Leu Cys Gly Ser Arg Lys
 195 200 205

Ile Pro Leu Thr Arg Leu Tyr Val Thr Ile Leu Leu Thr Val Leu Val
 30 210 215 220

SUBSTITUTE SHEET (RULE 26)

Phe Leu Leu Cys Gly Leu Pro Phe Gly Ile Gln Phe Phe Leu Phe Leu
 225 230 235 240

5 Trp Ile His Val Asp Arg Glu Val Leu Phe Cys His Val His Leu Val
 245 250 255

Ser Ile Phe Leu Ser Ala Leu Asn Ser Ser Ala Asn Pro Ile Ile Tyr
 260 265 270

10 Phe Phe Val Gly Ser Phe Arg Gln Arg Gln Asn Arg Gln Asn Leu Lys
 275 280 285

Leu Val Leu Gln Arg Ala Leu Gln Asp Ala Ser Glu Val Asp Glu Gly
 15 290 295 300

Gly Gly Gln Leu Pro Gln Glu Thr Leu Glu Leu Ser Gly Ser Arg Leu
 305 310 315 320

20 Glu Gln

(2) INFORMATION FOR SEQ ID NO:10:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 969 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

30

SUBSTITUTE SHEET (RULE 26)

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

5 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

ATGGATCCAA CGGTCTCAAC CTTGGACACA GAATTGACAC CAATCAACGG AACTGAGGAG 60
10
ACTCTTTGCT ACAAGCAGAC CTTGAGCCTC ACGGTGCTGA CGTGCATCGT TTCCCTTGTC 120
GGGCTGACAG GAAACGCGGT TGTGCTCTGG CTCCTGGGCT GCCGCATGCG CAGGAACGCC 180
15 TTCTCCATCT ACATCCTCAA CTTGGCCGCA GCAGACTTCC TCTCCTCAG CGGCCGCCTT 240
ATATATTCCC TGTAAAGCTT CATCAGTATC CCCCATACCA TCTCTAAAAT CCTCTATCCT 300
GTGATGATGT TTTCTACTT TGCAGGCCTG AGCTTTCTGA GTGCCGTGAG CACCGAGCGC 360
20
TGCCTGTCCG TCCTGTGGCC CATCTGGTAC CGCTGCCACC GCCCCACACA CCTGTCAGCG 420
GTGGTGTGTG TCCTGCTCTG GGCCCTGTCC CTGCTGCGGA GCATCCTGGA GTGGATGTTA 480
25 TGTGGCTTCC TGTCAGTGG TGCTGATTCT GCTTGGTGTC AAACATCAGA TTTCATCACA 540
GTCCGCTGGC TGATTTTTTT ATGTGTGGTT CTCTGTGGGT CCAGCCTGGT CCTGCTGATC 600
AGGATTCTCT GTGGATCCCG GAAGATACCG CTGACCAGGC TGTACGTGAC CATCCTGCTC 660
30

SUBSTITUTE SHEET (RULE 26)

ACAGTACTGG TCTTCCTCCT CTGTGGGCTG CCGTTTGGCA TTCAGTTTTT CCTATTTTTA 720
TGGATCCACG TGGACAGGGA AGTCTTATTT TGTCACTGTC ATCTAGTTTC TATTTTCCTG 780
5 TCCGCTCTTA ACAGCACTGC CAACCCCATC ATTTACTTCT TCGTGGGCTC CTTTAGGCAG 840
CGTCAAAATA GGCAGAACCT GAAGCTGGTT CTCCAGAGGG CTCTGCAGGA CGCGTCTGAG 900
GTGGATGAAG GTGGAGGGCA GCTTCCTCAG GAAATCCTGG AGCTGTCGGG AAGCAGATTG 960
10 GAGCAGTGA 969

(2) INFORMATION FOR SEQ ID NO:11:

15

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 322 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
20 (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

25

(iv) ANTI-SENSE: NO

Met Asp Pro Thr Val Pro Val Leu Gly Thr Lys Leu Thr Pro Ile Asn
5 1 5 10 15

Gly Arg Glu Glu Thr Pro Cys Tyr Lys Gln Thr Leu Ser Phe Thr Val
20 25 30

10 Leu Thr Cys Ile Ile Ser Leu Val Gly Leu Thr Gly Asn Ala Val Val
 35 40 45

Leu Trp Leu Leu Gly Cys Arg Met Arg Arg Asn Ala Val Ser Ile Tyr
50 55 60

15
Ile Leu Asn Leu Ala Ala Ala Asp Phe Leu Phe Leu Ser Phe Gln Ile
65 70 75 80

Ile Cys Arg Pro Leu Arg Leu Ile Asn Ile Ser His Leu Ile Arg Lys

Ile Leu Val Ser Val Met Thr Phe Pro Tyr Phe Thr Gly Leu Ser Met
100 105 110

25 Leu Ser Ala Ile Ser Thr Glu Arg Cys Leu Ser Val Leu Trp Pro Ile
 115 120 125

Trp Tyr Arg Cys Arg Arg Pro Thr His Leu Ser Ala Val Val Cys Val
130 135 140

30

BNSDOCID: <WO_9832519A1_1A>

Leu Leu Trp Ala Gly Leu Leu Leu Phe Ser Met Leu Glu Trp Arg Phe
 145 150 155 160

Cys Asp Phe Leu Phe Ser Gly Ala Asp Ser Ser Trp Cys Glu Thr Ser
 5 165 170 175

Asp Phe Ile Pro Val Ala Trp Leu Ile Phe Leu Cys Val Val Leu Cys
 180 185 190

Val Ser Ser Leu Val Leu Leu Val Arg Ile Leu Cys Gly Ser Arg Lys
 10 195 200 205

Met Pro Leu Thr Arg Leu Tyr Val Thr Ile Leu Leu Thr Val Leu Val
 210 215 220

Phe Leu Leu Cys Gly Leu Pro Phe Gly Ile Leu Gly Ala Leu Ile Tyr
 15 225 230 235 240

Arg Met His Leu Asn Leu Glu Val Leu Tyr Cys His Val Tyr Leu Val
 20 245 250 255

Cys Met Ser Leu Ser Ser Leu Asn Ser Ser Ala Asn Pro Ile Ile Tyr
 260 265 270

Phe Phe Val Gly Ser Phe Arg Gln Arg Gln Asn Arg Gln Asn Leu Lys
 25 275 280 285

Leu Val Leu Gln Arg Ala Leu Gln Asp Lys Pro Glu Val Asp Lys Gly
 290 295 300

30

SUBSTITUTE SHEET (RULE 26)

Glu Gly Gln Leu Pro Glu Glu Ser Leu Glu Leu Ser Gly Arg Arg Leu

305

310

315

320

Gly Pro

5

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

10

(A) LENGTH: 969 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

ATGGATCCAA CCGTCCCAGT CTTGGGTACA AAACGACAC CAATCAACGG ACGTGAGGAG 60

15 ACTCCTTGCT ACAAGCAGAC CCTGAGCTTC ACGGTGCTGA CGTGATCAT TTCCCTTGTC 120

GGACTGACAG GAAACGCGGT TGTGCTCTGG CTCCTGGGCT GCCGCATGCG CAGGAACGCT 180

GTCTCCATCT ACATCCTCAA CCTGGCCGCA GCAGACTTCC TCTTCCTCAG CTTCCAAATT 240

30

SUBSTITUTE SHEET (RULE 26)

ATACGTTCCG CATTACGGCT CATCAATATC AGCCATCTCA TCCGCAAAAT CCTCGTTTCT 300

GTGATGACCT TTCCCTACTT TACAGGGCTG AGTATGCTGA GCSCCATCAG CACCGAGCGC 360

5 TGCCTGTCTG TTCTGTGGCC CATCTGGTAC CGCTGCCGCC GCCCCACACA CCTGTCAGCG 420

GTCTGTGTG TCCTGCTCTG GGGCCTGTCC CTGCTGTTTA GTATGCTGGA GTGGAGGTTT 480

TGTGACTTCC TGTTTAGTGG TGCTGATTCT AGTTGGTGTG AAACGTCAGA TTTCATCCCA 540

10 GTCCCGTGGC TGATTTTTTT ATGTGTGGTT CTCTGTGTTT CCAGCCTGGT CCTGCTGGTC 600

AGGATCCTCT GTGGATCCCG GAAGATGCCG CTGACCAGGC TGTATGTGAC CATCCTGCTC 660

15 ACAGTGCTGG TCTTCCTCCT CTGCGGCCCTG CCTTCGGCA TTCTGGGGGC CCTAATTAC 720

AGGATGCACC TGAATTTGGA AGTCTTATAT TGTATGTTT ATCTGGTTTG CATGTCCCTG 780

TCCTCTCTAA ACAGTAGTGC CAACCCCATC ATTTACTTCT TCGTGGGCTC CTTTAGGCAG 840

20 CGTCAAAATA GGCAGAACCT GAAGCTGGTT CTCCAGAGGG CTCTGCAGGA CAAGCCTGAG 900

GTGGATAAAG GTGAAGGGCA GCTTCCTGAG GAAAGCCTGG AGCTGTCGGG AAGGAGATTG 960

25 GGGCCATGA 969

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 322 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: not relevant

10 (ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Met Asp Pro Thr Val Pro Val Phe Gly Thr Lys Leu Thr Pro Ile Asn

1 5 10 15

20

Gly Arg Glu Glu Thr Pro Cys Tyr Asn Gln Thr Leu Ser Phe Thr Val

20 25 30

Leu Thr Cys Ile Ile Ser Leu Val Gly Leu Thr Gly Asn Ala Val Val

25 35 40 45

Leu Trp Leu Leu Gly Tyr Arg Met Arg Arg Asn Ala Val Ser Ile Tyr

50 55 60

SUBSTITUTE SHEET (RULE 26)

Ile Leu Asn Leu Ala Ala Ala Asp Phe Leu Phe Leu Ser Phe Gln Ile
 65 70 75 80

5 Ile Arg Ser Pro Leu Arg Leu Ile Asn Ile Ser His Leu Ile Arg Lys
 85 90 95

Ile Leu Val Ser Val Met Thr Phe Pro Tyr Phe Thr Gly Leu Ser Met
 100 105 110

10 Leu Ser Ala Ile Ser Thr Glu Arg Cys Leu Ser Val Leu Trp Pro Ile
 115 120 125

Trp Tyr Arg Cys Arg Arg Pro Thr His Leu Ser Ala Val Val Cys Val
 15 130 135 140

Leu Leu Trp Gly Leu Ser Leu Leu Phe Ser Met Leu Glu Trp Arg Phe
 145 150 155 160

20 Cys Asp Phe Leu Phe Ser Gly Ala Asp Ser Ser Trp Cys Glu Thr Ser
 165 170 175

Asp Phe Ile Pro Val Val Trp Leu Ile Phe Leu Cys Val Val Leu Cys
 180 185 190

25 Val Ser Ser Leu Val Leu Leu Val Arg Ile Leu Cys Gly Ser Arg Lys
 195 200 205

Met Pro Leu Thr Arg Leu Tyr Val Thr Ile Leu Leu Thr Val Leu Val
 30 210 215 220

SUBSTITUTE SHEET (RULE 26)

Phe Leu Leu Cys Gly Leu Pro Phe Gly Ile Leu Gly Ala Leu Ile Tyr
 225 230 235 240

5 Arg Met His Leu Asn Leu Glu Val Leu Tyr Cys His Val Tyr Leu Val
 245 250 255

Cys Met Ser Leu Ser Ser Leu Asn Ser Ser Ala Asn Pro Ile Ile Tyr
 260 265 270

10 Phe Phe Val Gly Ser Phe Arg Gln Arg Gln Asn Arg Gln Asn Leu Lys
 275 280 285

Leu Val Leu Gln Arg Ala Leu Gln Asp Lys Pro Glu Val Asp Lys Gly
 15 290 295 300

Glu Gly Gln Leu Pro Glu Glu Ser Leu Glu Leu Ser Gly Ser Lys Leu
 305 310 315 320

20 Gly Pro

(2) INFORMATION FOR SEQ ID NO:14:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 969 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

30

SUBSTITUTE SHEET (RULE 26)

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

5 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

ATGGATCCAA CCGTCCCAGT CTTCGGTACA AAAGTGACAC CAATCAACGG ACGTGAGGAG 60
10 ACTCCTTGCT ACAATCAGAC CCTGAGCTTC ACGGTGCTGA CGTGATCAT TTCCCTTGTC 120
GGACTGACAG GAAACGCGGT TGTGCTCTGG CTCCTGGGCT ACCGCATGCG CAGGAACGCT 180
15 GTCTCCATCT ACATCCTCAA CCTGGCCGCA GCAGACTTCC TCTTCTCAG CTTCCAAATT 240
ATACGTTGCG CATTACGCCT CATCAATATC AGCCATCTCA TCCGCAAAAT CCTCGTTTCT 300
GTGATGACCT TTCCCTACTT TACAGGCCTG AGTATGCTGA GCGCCATCAG CACCGAGCGC 360
20 TGCCTGTCTG TTCTGTGGCC CATCTGGTAC CGCTGCCGCC GCCCCACACA CCTGTCAGCG 420
GTCGTGTGTG TCCTGCTCTG GGGCCTGTCC CTGCTGTTTA GTATGCTGGA GTGGAGGTTT 480
25 TGTGACTTCC TGTTTAGTGG TGCTGATTCT AGTTGGTGTG AAACGTCAGA TTTCATCCCA 540
GTGCTGTGGC TGATTTTTTT ATGTGTGGTT CTCTGTSTTT CCAGCCTGGT CCTGCTGGTC 600
AGGATCCTCT GTGGATCCCG GAAGATGCCG CTGACCAGGC TGTACGTGAC CATCCTGCTC 660
30

SUBSTITUTE SHEET (RULE 26)

ACAGTGCTGG TCTTCTCTCT CTGCGGCGCTG CCCTTCGGCA TTCTGGGGGC CCTAATTTAC 720

AGGATGCACC TGAATTTGGA AGTCTTATAT TGTCTGTTT ATCTGGTTTG CATGTCCCTG 780

5 TCCTCTCTAA ACAGTAGTGC CAACCCCATC ATTTACTTCT TCGTGGGCTC CTTTAGGCAG 840

CGTCAAAATA GGCAGAACCT GAAGCTGGTT CTCCAAAGGG CTCTGCAGGA CAAGCCTGAG 900

GTGGATAAAG GTGAAGGGCA GCTTCTGAG GAAAGCCTGG AGCTGTCGGG AAGCAAATTG 960

10 GGGCCATGA 969

(2) INFORMATION FOR SEQ ID NO:15:

15

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

20

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic PCR primer"

25

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

30

SUBSTITUTE SHEET (RULE 26)

GGCCGTCGAC TTCATCGTCW MYCTIKCIYT IGCNG

35

(2) INFORMATION FOR SEQ ID NO:16:

5

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

10

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic PCR primer"

15

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

20

RHWRCARTAI ATIAAT

15

(2) INFORMATION FOR SEQ ID NO:17:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

30

(D) TOPOLOGY: linear

SUBSTITUTE SHEET (RULE 26)

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic PCR primer"

5 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

10

CGCAGATGAG GTAGTACAGC ATCAC

25

(2) INFORMATION FOR SEQ ID NO:18:

15

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

20 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic PCR primer"

25 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

30

SUBSTITUTE SHEET (RULE 26)

CTGTGAGAGA GATGGTAACA TACAG

25

(2) INFORMATION FOR SEQ ID NO:19:

5

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

10

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic PCR primer"

15

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

20

GCATCCTTGA CTGGTCTTC TCAG

24

(2) INFORMATION FOR SEQ ID NO:20:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

30

(D) TOPOLOGY: linear

SUBSTITUTE SHEET (RULE 26)

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic PCR primer"

5 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

10

GGGTGAGACT CATCATCATT TGTGG

25

(2) INFORMATION FOR SEQ ID NO:21:

15

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

20

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic PCR primer"

(iii) HYPOTHETICAL: NO

25

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

30 GCAAGCTTTC TGAGCATGGA TCCAACCGTC

30

SUBSTITUTE SHEET (RULE 26)

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "synthetic PCR primer"

(iii) HYPOTHETICAL: NO

15 (iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

CCCTCAGATC TCCAATTTCG TTCCCGACAG

30

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 98/02348

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07K 14/72

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9504073 A1 (THE BOARD OF TRUSTEES OF THE LELAND STANFORD JUNIOR UNIVERSITY), 9 February 1995 (09.02.95)	1-35, 38-42
A	--	36-37
X	WO 9405695 A1 (NEW YORK UNIVERSITY), 17 March 1994 (17.03.94), See page 14, line 11 seq 52, claims	1-42
A	Dialog Information Service, file 154, Medline, Dialog accession no. 08044093, Medline accession no. 95047685, Brown NJ et al: "Gastrointestinal adaptation to enhanced small intestinal lipid exposure", Gut (ENGLAND) Oct 1994, 35 (10) p 1409-12	1-42
	--	

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

6 April 1999

Date of mailing of the international search report

14.04.99

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Patrick Andersson

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

Information on patent family members

02/03/99

International application No.

PCT/SE 98/02348

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9504073	A1	09/02/95	US	5556780 A	17/09/96
WO	9405695	A1	17/03/94	AU	4855393 A	29/03/94
				US	5508384 A	16/04/96